

REFERÊNCIAS BIBLIOGRÁFICAS

Aptamil Pepti – a fórmula com nutrientes mais próximos do padrão ouro no manejo da APLV

1. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am.* 2013;60(1):49-74.
2. Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. *Arq Asma Alerg Imunol.* 2018;2(1):7-82.
3. Venter C et al. Diagnosis and management of non-IgE-mediated cow's milk allergy in infancy: a UK primary care practical guide. *Clin Transl Allergy.* 2013;3(1):23
4. Vandenplas Y, Broekaert I, Domellöf M, Indrio F, Lapillonne A, Pienar C, Ribes-Koninckx C, Shamir R, Szajewska H, Thapar N, Thomassen RA, Verduci E, West C. An ESPGHAN Position Paper on the Diagnosis, Management, and Prevention of Cow's Milk Allergy. *J Pediatr Gastroenterol Nutr.* 2024 Feb;78(2):386-413.
5. Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.
6. gg
7. Vandenplas Y. Lactose Intolerance. *Asia Pac J Clin Nutr.* 2015;24(Suppl 1):S9-13
8. Arslanoglu S, Moro GE, Boehm G, Wienz F, Stahl B, Bertino E. Early neutral prebiotic oligosaccharide supplementation reduces the incidence of some allergic manifestations in the first 5 years of life. *J Biol Regul Homeost Agents.* 2012;26(3 Suppl):49-59.
9. Arslanoglu S, Moro GE, Schmitt J, Tandoi L, Rizzardi S, Boehm G. Early Dietary intervention with a mixture of prebiotic oligosaccharides reduces the incidence of allergic manifestations and infections during the first two years of life. *J Nutr.* 2008;138(6):1091-5.
10. Moro G, Arslanoglu S, Stahl B, Jelinek J, Wahn U, 12. Boehm G. A mixture of prebiotics oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Arch Dis Child.* 2006;91(10):814-9.
11. Koletzko B, Baker S, Cleghorn G, et al. Global standard for the composition of infant formula: recommendations of an ESPGHAN coordinated international expert group. *Pediatr Gastroenterol Nutr.* 2005;41(5):584-99.
12. Birch E. et al. The DIAMOND (DHA intake and measurement of neural development) study: a double-masked, randomized controlled clinical trial of the maturation of infant visual acuity as a function of the dietary level of docosahexaenoic acid' *Am J Clin Nutr.*, vol. 91, no.4, pg. 848-859, 2010.
13. Qawasmi A et al. Meta-analysis of LCPUFA supplementation of infant formula and visual acuity. *Pediatrics.* 2013;131(1):262-72.
14. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
15. Miles EA, Childs CE, Calder PC. Long-Chain Polyunsaturated Fatty Acids (LCPUFAs) and the Developing Immune System: A Narrative Review. *Nutrients.* 2021; 13: 247

16. Estudo observacional, prospectivo, aberto e multicêntrico de vigilância pós-comercialização sobre aceitabilidade e tolerabilidade de uma fórmula infantil extensamente hidrolisada. Brasil. 2016: dados preliminares.
17. Epifanio, M. et al. Palatability and Sensory Perception of Infant Formulas for the Treatment of Cow's Milk Allergy According to Brazilian Mothers. *Biomed J Sci & Tech Res* 25(1)-2020. BJSTR. MS.ID.00416.
18. IQVIA, 2023
19. Verwimp JJ, et al. Symptomatology and growth in infants with cow's milk protein intolerance using two different whey protein hydrolysate based formulas in a Primary Health Care setting. *Eur J Clin Nutr.* 1995;49 Suppl 1:S39 S48.2.
20. Giampietro PG, et al. Hypoallergenicity of an extensively hydrolyzed whey formula. *Pediatr Allergy Immunol.* 2001;12(2):83 86. 3.
21. Isolauri, E., et al (1995). Efficacy and safety of hydrolyzed cow milk and amino acid derived formulas in infants with cow milk allergy. *The Journal of pediatrics*, 127(4), 550 557.
22. Sampson et al (1992). Safety of an amino acid derived infant formula in children allergic to cow milk. *Pediatrics*, 90(3), 463 465.4.
23. Meyer R, Venter C, Fox AT, Shah N. Practical dietary management of protein energy malnutrition in young children with cow's milk protein allergy. *Pediatr Allergy Immunol.* 2012;23(4):307–14.
24. Moraes MB, Spolidoro JV, Vieira MC, Cardoso AL, Clark O, Nishikawa A, Castro AP. Amino acid formula as a new strategy for diagnosing cow's milk allergy in infants: is it cost-effective? *J Med Econ.* 2016;19:1207-14.
25. Meyer R et al. When Should Infants with Cow's Milk Protein Allergy Use an Amino Acid Formula? A Practical Guide. *J Allergy Clin Immunol Pract.* 2018 Mar - Apr;6(2):383-399.
26. Muraro A et al. EAACI food allergy and anaphylaxis guidelines: diagnosis and management of food allergy. *Allergy.* 2014;69(8):1008-25.