

REFERÊNCIAS BIBLIOGRÁFICAS

Monografia Aptamil Pepti HMO

Referências Capa

1. de Oliveira LCL, et al. Arq Asma Alerg Imunol – Vol. 9, Nº 1, 2025
2. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. Pediatr Clin North Am.2013;60(1):49-74.
3. Vandenplas Y, et al. An ESPGHAN Position Paper on the Diagnosis, Management, and Prevention of Cow's Milk Allergy. J Pediatr Gastroenterol Nutr. 2024 Feb;78(2):386-413.
4. Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.

Referências telas de Produto

5. de Oliveira LCL, et al. Arq Asma Alerg Imunol – Vol. 9, Nº 1, 2025
6. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. Pediatr Clin North Am.2013;60(1):49-74.
7. Vandenplas Y, et al. An ESPGHAN Position Paper on the Diagnosis, Management, and Prevention of Cow's Milk Allergy. J Pediatr Gastroenterol Nutr. 2024 Feb;78(2):386-413.
8. Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.
9. Nutrologia Pediátrica: Temas da Atualidade em Nutrologia Pediátrica - 2021. / Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia. São Paulo: SBP, 2021.
10. FERREIRA, CHT; BARRETO, BAP; TOFOLI, MHC; et al. Microbioma e pediatria na saúde e na doença. Barueri [SP]: Manole, 2024. 1.ed.
11. Arslanoglu S, Moro GE, Boehm G, Wenz F, Stahl B, Bertino E. Early neutral prebiotic oligosaccharide supplementation reduces the incidence of some allergic manifestations in the first 5 years of life. J Biol Regul Homeost Agents. 2012;26(3 Suppl):49-59.
12. Arslanoglu S, Moro GE, Schmitt J, Tandoi L, Rizzardi S, Boehm G. Early Dietary intervention with a mixture of prebiotic oligosaccharides reduces the incidence of allergic manifestations and infections during the first two years of life. J Nutr. 2008;138(6):1091-5.
13. Boehm G. A mixture of prebiotics oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. Arch Dis Child. 2006;91(10):814-

14. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5
15. Koletzko B, Baker S, Cleghorn G, et al. Global standard for the composition of infant formula: recommendations of an ESPGHAN coordinated international expert group. *J Pediatr Gastroenterol Nutr.* 2005;41(5):584-99.
16. SALMINEN, Seppo; SZAJEWSKA, Hania; KNOL, Jan. *The Biotics Family in Early Life*. 2. ed. Wiley, 2023.
17. Goh CY, et al. Combination of short-chain GOS and long-chain FOS 9:1 with 2FL positively impact the infant gut microbiota composition and metabolic activity in a stimulator of the human intestinal microbial ecosystem (SHIME). *J Pediatc Gastroenterol Nutr* 2019; 68 (S1): NP 114; 1159.
18. Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar/ Sociedade Brasileira de Pediatria. – 5.ed. rev. ampl.– São Paulo: SBP, 2024. 208 p.
19. Bode L. Human milk oligosaccharides: every baby needs a sugar mama. *Glycobiology.* 2012 Sep;22(9):1147-62.
20. Estudo observacional, prospectivo, aberto e multicêntrico de vigilância pós-comercialização sobre aceitabilidade e tolerabilidade de uma fórmula infantil extensamente hidrolisada. Brasil. 2016: dados preliminares.
21. Epifanio, M. et al. Palatability and Sensory Perception of Infant Formulas for the Treatment of Cow's Milk Allergy According to Brazilian Mothers. *Biomed J Sci & Tech Res* 25(1)-2020. BJSTR. MS.ID.00416.
22. Birch E. et al. The DIAMOND (DHA intake and measurement of neural development) study: a double-masked, randomized controlled clinical trial of the maturation of infant visual acuity as a function of the dietary level of docosahexaenoic acid' *Am J Clin Nutr.*, vol. 91, no.4, pg. 848–859, 2010.
23. Qawasmi A et al. Meta-analysis of LCPUFA supplementation of infant formula and visual acuity. *Pediatrics.* 2013;131(1):262-72.
24. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
25. Miles EA, Childs CE, Calder PC. Long-Chain Polyunsaturated Fatty Acids (LCPUFAs) and the Developing Immune System: A Narrative Review. *Nutrients.* 2021; 13: 247
26. HCP TRACKER, 2024
27. Verwimp JJ, et al. Symptomatology and growth in infants with cow's milk protein intolerance using two different whey protein hydrolysate based formulas in a Primary Health Care setting. *Eur J Clin Nutr.* 1995;49 Suppl 1:S39 S48.2.
28. Giampietro PG, et al. Hypoallergenicity of an extensively hydrolyzed whey formula. *Pediatr Allergy Immunol.* 2001;12(2):83 86. 3.
29. Sampson et al (1992). Safety of an amino acid derived infant formula in children allergic to cow milk. *Pediatrics,* 90(3), 463 465.4.

30. Isolauri, E., et al (1995). Efficacy and safety of hydrolyzed cow milk and amino acid derived formulas in infants with cow milk allergy. *The Journal of pediatrics*, 127(4), 550-557.
31. Morais MB, Spolidoro JV, Vieira MC, Cardoso AL, Clark O, Nishikawa A, Castro AP. Amino acid formula as a new strategy for diagnosing cow's milk allergy in infants: is it cost-effective? *J Med Econ.* 2016;19:1207-14.
32. Meyer R et al. When Should Infants with Cow's Milk Protein Allergy Use an Amino Acid Formula? A Practical Guide. *J Allergy Clin Immunol Pract.* 2018 Mar - Apr;6(2):383-399.
33. SBP. Guia de orientações - Dificuldades alimentares/ Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia . São Paulo: SBP, 2022. 66
34. Meyer R, Venter C, Fox AT, Shah N. Practical dietary management of protein energy malnutrition in young children with cow's milk protein allergy. *Pediatr Allergy Immunol.* 2012;23(4):307-14.
35. Eveleens, R D et al. Definitions, predictors, and outcomes of feeding intolerance in critically ill children: A systematic review. *Clin Nutr,* 2020, 39.3: 685-693.
36. Marino, LV et al. Feeding intolerance in children with critical illness. *Clin. Nutr.,* 2020, 39.3: 609-611.
37. Lucia Diaferio L et al. May Failure to Thrive in Infants Be a Clinical Marker for the Early Diagnosis of Cow's Milk Allergy?. *Nutrients .* 2020 Feb 13;12(2):466. doi: 10.3390/nu12020466.