

Linha Aptamil Pre e Pre Transition

REFERÊNCIAS BIBLIOGRÁFICAS

- 1.Koletzko, Berthold, et al. “Scientific basis and practical application of nutritional care for preterm infants.” Nutritional Care of Preterm Infants. Vol. 122. Karger Publishers, 2021.
- 2.Embleton, N. D., Moltu, S. J., Lapillonne et al. Enteral nutrition in preterm infants (2022): a position paper from the ESPGHAN committee on nutrition and invited experts. Journal of pediatric gastroenterology and nutrition.2023; 76(2), 248-268.
- 3.Aggett, P. J. Agostoni, C. Axelsson I. et al. 2006 Feeding preterm infants after hospital discharge: a commentary by the ESPGHAN Committee on Nutrition. Journal of Pediatric Gastroenterology and Nutrition, 42, 596-603.
- 4.Roy CC et al. Correction of the malabsorption of the preterm infant with a medium-chain triglyceride formula. J Pediatr. 1975;86(3):446-50.
- 5.Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. BRASPEN J 2020; 35 (3): 294-306.
- 6.Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. J Pediatr Gastroenterol Nutr. 2002;34(3):291-5.
- 7.Knol J et al. Increase of faecal bifidobacteria due to dietary oligosaccharides induces a reduction of clinically relevant pathogen germs in the faeces of formula-fed preterm infants. Acta Pædiatrica, 2005; 94(Suppl 449): 31–33.
- 8.Bruzzese E et al. Early administration of GOS/FOS prevents intestinal and respiratory infections in infants. J Pediatr Gastroenterol Nutr. 2006; 42 (5): E95.
- 9.Westerbeek EAM et al. Neutral and acidic oligosaccharides in preterm infants: a randomized, double-blind, placebo-controlled trial. Am J Clin Nutr 2010; 91:679-86.
- 10.Westerbeek EAM et al. Design of a randomised controlled trial on immune effects of acidic and neutral oligosaccharides in the nutrition of preterm infants: carrot study. BMC Pediatrics 2008, 8:46.
- 11.Liu L et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. J Lipid Res. 2014;55(3):531-9.
- 12.Bar-Yoseph F et al. Review of sn-2 palmitate oil implications for infant health. Prostaglandins Leukot Essent Fatty Acids. 2013;89(4):139-43.
- 13.Yaron, S. et al. Nutr. 2013 Apr;56(4):376-81 Effect of high-palmitate content in infant formula on the intestinal microbiota of term infants. J Pediatr Gastroenterol.
- 14.Comparativo realizado entre produtos da mesma categoria, em 08/2025, utilizando informações extraídas da tabela nutricional.

A importância dos lipídios na nutrição do prematuro

REFERÊNCIAS BIBLIOGRÁFICAS

1. Aggett, P.J. et al (2006). Feeding preterm infants after hospital discharge: a commentary by the ESPGHAN Committee on Nutrition. *Journal of Pediatric Gastroenterology and Nutrition*, 42(5), 596–603.
2. Embleto, N. D. (2023). Enteral nutrition in preterm infants (2022): a position paper from the ESPGHAN Committee on Nutrition and Invited Experts. *Journal of Pediatric Gastroenterology and Nutrition*, 76 (2), 248–268.
3. Falcão, M.C. (2020). Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN Journal*, 35(3), 294–306.
4. Havlicekova, S. et al (2016). Beta-palmitate – a natural component of human milk in supplemental milk formulas. *Nutrition Journal*, 15(28), 1–8.
5. Hellström, A. et al (2021). Effect of enteral lipid supplement on severe retinopathy of prematurity - a Randomized Clinical Trial. *JAMA Pediatrics*, 175(4), 359–367.
6. Hu, R. et al (2024). Could early life DHA supplementation benefit neurodevelopment? A systematic review and metaanalysis. *Frontiers in Neurology*, 15, 01–17.
7. Martini, C. A. (2006). Ácidos graxos poliinsaturados ômega-3 e ômega-6: importância e ocorrência em alimentos. *Revista de Nutrição*, 19 (6), 761–770.
8. Moltu, S. J. et al (2024). Arachidonic and docosahexaenoic acid supplementation and brain maturation in preterm infants; a double-blind RCT. *Clinical Nutrition*, 43, 176–186.
9. Nowacki, J. et al (2014). Stool fatty acid soaps, stool consistency and gastrointestinal tolerance in term infants fed infant formulas containing high sn-2 palmitate with or without oligofructose: a double-blind, randomized clinical trial. *Nutrition Journal*, 5(13), 105–115.
10. Quintan, P.T. et al (1995). The relationship between stool hardness and stool composition in breast- and formula-fed infants. *Journal of Pediatric Gastroenterology and Nutrition*, 20(1), 81–90.
11. SBP (2024). Manual de seguimento do recém-nascido de alto risco/ Departamento Científico de Neonatologia da Sociedade Brasileira de Pediatria. - 2.ed.- São Paulo, 142 f.
12. Silva, D.R.B. et al (2007). A importância dos ácidos graxos poliinsaturados de cadeia longa na gestação e lactação. *Revista Brasileira de Saúde Materno Infantil*, 7 (2), 123–133.
13. Vanek, V.W. et al. (2012). A.S.P.E.N. position paper: clinical role for alternative intravenous fat emulsions. *Nutrition in Clinical Practice*, 27(2), 150–192.
14. Wendel, K. et al (2023). Essential fatty acid supplementation and early inflammation in preterm infants: secondary analysis of a randomized clinical trial. *Neonatology*, 120, 465–472.

15. Yao, M. et al (2014). Effects of term infant formulas containing high sn-2 palmitate with and without oligofructose on stool composition, stool characteristics, and bifidogenicity. *Journal of Pediatric Gastroenterology and Nutrition*, 59(4), 440–448.
16. Yuan, T. et al (2022). Role medium-chain fatty acids in the lipid metabolism of infants. *Frontiers in Nutrition*, 9, 1–7.