

REFERÊNCIAS BIBLIOGRÁFICAS

Mitos e Verdades – Infatrini

1. ASPEN. Nutrition Management of Term Infants with Growth Failure. 2022
2. Codex Alimentarius FAO/OMS. Standard for infant formula and formulas for special medical purposes intended for infants. Codex Stan 72 – 1981. Revised 2007/2011.
3. Gahagan S. Failure to Thrive: A Consequence of Undernutrition. Pediatrics in Review Vol.27 No.1 January 2006
4. Clarke SE, Evans S, MacDonald A, Davies P, Booth IW. Randomized comparison of a nutrient-dense formula with an energy-supplemented formula for infants with faltering growth. J Hum Nutr Diet. 2007;20(4):329-339.
5. Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4ª ed. Rio de Janeiro, 2018.
6. Braegger C, Decsi T, Dias JA, Hartman C, Kolaček S, Koletzko B et al. Practical Approach to Paediatric Enteral Nutrition: A Comment by the ESPGHAN Committee on Nutrition. JPGN; 2010;51(1):110-122
7. Manual de Suporte Nutricional da Sociedade Brasileira de Pediatria - 2ed. - Rio de Janeiro: Departamento Científico de Suporte Nutricional da Sociedade Brasileira de Pediatria. – 2020.
8. Antonella, et al. “Prevalence and outcome of malnutrition in pediatric patients with chronic diseases: Focus on the settings of care.” Clinical Nutrition 38.4 (2019): 1877-1882.
9. Joosten K, Meyer R. Nutritional screening and guidelines for managing the child with faltering growth. EJCN. 2010;64:S23-S24
10. Habibzadeh, H. Jafarizadeh, A. DIDARLOO. Determinants of failure to thrive (FTT) among infants aged 6-24 months: a case-control study. J prev med hyg 2015; 56: E180-E186.
11. World Health Organization. Safe preparation, storage and handling of powdered infant
12. formula [Online] 2007. Available at:
http://www.who.int/foodsafety/publications/micro/pif_guidelines.pdf
13. World Health Organization. Protein and amino acid requirements in human nutrition. Report of a Joint WHO/FAO/UNU Expert Consultation. World Health Organ Tech Rep Ser 2007; (935): 1– 265
14. Steele JR, et al. Determining the osmolality of over-concentrated and supplemented infant formulas. J Hum Nutr Diet. 2013 Feb;26(1):32-7.
15. Slicker J, et al. Nutrition algorithms for infants with hypoplastic left heart

syndrome; birth through the first interstage period. *Congenit Heart Dis*. 2013 Mar-Apr;8(2):89-102.

16. Roman, B. Nourishing little hearts: Nutritional implications for congenital heart defects. *Practical Gastroenterology*. 2011.35(8):11-34.

17. Pereira-da-Silva L, et al. Osmolality of preterm formulas supplemented with nonprotein energy supplements. *Eur J Clin Nutr*. 2008 Feb;62(2):274-8.

18. Alsoufi B, et al. Low-weight infants are at increased mortality risk after palliative or corrective cardiac surgery. *J Thorac Cardiovasc Surg*. 2014 Dec;148(6):2508-14.e1.

19. Reddy VM. Low birth weight and very low birth weight neonates with congenital heart disease: timing of surgery, reasons for delaying or not delaying surgery. *Semin Thorac Cardiovasc Surg Pediatr Card Surg Annu*. 2013;16(1):13-20.

20. Fomon SJ, Ziegler EE. Renal solute load and potential renal solute load in infancy. *J Pediatr*. 1999 Jan;134(1):11-4.

21. Carvalho LR, et al. Hazard analysis and critical control point system approach in the evaluation of environmental and procedural sources of contamination of enteral feedings in three hospitals. *JPEN J Parenter Enteral Nutr*. 2000 Sep-Oct;24(5):296-303.

22. Fagerman KE. Limiting bacterial contamination of enteral nutrient solutions: 6-year history with reduction of contamination at two institutions. *Nutr Clin Pract*. 1992 Feb;7(1):31-6.

23. Labiner-Wolfe J, et al. Infant formula-handling education and safety. *Pediatrics*. 2008 Oct;122 Suppl 2:S85-90.

24. Pencharz PB. Protein and energy requirements for “optimal” catch-up growth. *EJCN*.2010;64: S5-7.

25. Mehta NM et al. Defining pediatric malnutrition: A paradigm shift toward etiology-related definitions. *J Parenter Enter Nutr*. 2013;37(4):460–81

26. Eveleens RD, Joosten KFM, de Koning BAE, Hulst JM, Verbruggen SCAT. Definitions, predictors and outcomes of feeding intolerance in critically ill children: A systematic review. *Clin Nutr*. 2020 Mar;39(3):685-693.

27. Marino LV, Johnson MJ, Davies NJ, Kidd CS, Fienberg J, Richens T, Bharucha T, Beattie RM, Darlington AE. Improving growth of infants with congenital heart disease using a consensus-based nutritional pathway. *Clin Nutr*. 2020 Aug;39(8).

28. Diaferio L, Caimmi D, Verga MC, Palladino V, Trovè L, Giordano P, Verduci E, Miniello VL. May Failure to Thrive in Infants Be a Clinical Marker for the Early Diagnosis of Cow's Milk Allergy? *Nutrients*. 2020 Feb 13;12(2):466. doi: 10.3390/nu12020466. PMID: 32069783; PMCID: PMC7071281.

29. Comparado a fórmula infantil para lactentes e de seguimento para lactentes e crianças de primeira infância. Setembro 2024.

30. FAO/WHO/UNU. Protein and amino acid requirements in human nutrition: report of a joint
31. FAO/WHO/UNU expert consultation. WHO technical report series no. 935, 2007.p.187
32. Falcão MC, Udsen N, Zamberlan P, Ceccon MER. Rev Bras Nutr Clin 2009; 24 (2): 125-30.
33. Uauy R, Mena P, Rojas C. Essential fatty acids in early life: structural and functional role. Proc Nutr Soc. 2000; 59:3-15.
34. Uauy R, Hoffman DR, Mena P, Llanos A, Birch EE. Term infant studies of DHA and ARA supplementation on neurodevelopment: results of randomized controlled trials. J Pediatr. 2003;143:S17-25.
35. SanGiovanni JP, Berkey CS, Dwyer JT, Colditz GA. Dietary essential fatty acids, long-chain polyunsaturated fatty acids, and visual resolution acuity in healthy full term infants: a systematic review. Early Hum Dev. 2000;57:165-88. 5
36. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. J Pediatr Gastroenterol Nutr. 2002;34(3):291-5.
37. Knol J et al. Colon microflora in infants fed formula with galacto- and fructo-oligosaccharides: more like breast-fed infants. J Pediatr Gastroenterol Nutr. 2005;
38. Bruzzese, Eugenia, et al. "A formula containing galacto-and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study." Clinical Nutrition 28.2 (2009): 156-161.