

REFERÊNCIAS

ONE PEDIATRIA DANONE

1. Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4ª ed. Rio de Janeiro, 2018.
2. Zubler et al. Evidence-informed milestones for developmental surveillance tools Pediatrics. 2022;149 (3)
3. Nogueira-de-Almeida, C. A., Falcão, M. C., Ribas Filho, D., Zorzo, R. A., Konstantyner, T., Ricci, R., Gioia, N., & Fisberg, M. (2022). Consensus of the Brazilian Association of Nutrology on Milky Feeding of Children Aged 1–5 Years Old. International Journal of Nutrology, 13(1), 2–16.
4. Manual de Suporte Nutricional da Sociedade Brasileira de Pediatria - 2ed. - Rio de Janeiro: Departamento Científico de Suporte Nutricional da Sociedade Brasileira de Pediatria. – 2020.
5. SBV. Pesquisa do IBOPE aponta crescimento histórico no número de vegetarianos no Brasil, Disponível em: Pesquisa do IBOPE aponta crescimento histórico no número de vegetarianos no Brasil - SVB. Acesso em 01.03.24.
6. Savilahti et al. 2012. 29.
7. Merritt, R. et al. North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition Position Paper: Plantbased Milks. JPN Volume 71, Number 2, August 2020.
8. Sociedade Brasileira de Pediatria. Guia de orientações - Dificuldades alimentares. Departamento Científico de Nutrologia São Paulo: SBP, 2022.
9. Codex Alimentarius FAO/OMS. Standard for infant formula and formulas for special medical purposes intended for infants. Codex Stan 72 – 1981. Revised 2007/2011.
10. Gahagan S. Failure to thrive: a consequence of undernutrition. Pediatr Rev. 2006 Jan;27(1):e1-11.
11. Benninga MA, Faure C, Hyman PE, et al. Childhood functional gastrointestinal disorders:neonate/toddler. Gastroenterology 2016; 150:1443-55.
12. Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation and cow's milk allergy in formula-fed infants. Acta Paediatrica 2015, ISSN 0803-5253
13. Savino F et al. Reduction of crying episodes owing to infantile colic: A randomized controlled study on the efficacy of a new infant formula. Eur J Clin Nutr 2006, 60:1304-10.
14. Savino F et al. "Minor" feeding problems during the first months of life: effect of a partially hydrolyzed milk formula containing fructo- and galacto-oligosaccharides. Acta Paediatr Suppl 2003, 91:86-90.
15. Schmelzle H et al. Randomized double-blind study of the nutritional efficacy and bifidogenicity of a new infant formula containing partially hydrolyzed protein, a high beta-palmitic acid level, and nondigestible oligosaccharides. J Pediatr Gastroenterol Nutr 2003, 36:343-51.
16. Bongers ME et al. The clinical effect of a new infant formula in term infants with constipation: a double-blind, randomized cross-over trial. Nutr J 2007; 6:8.
17. Savino F et al. Advances in the management of digestive problems during the first months of life. Acta Paediatr Suppl 2005, 94:120.
18. Borrelli O et al. Use of a new thickened formula for treatment of symptomatic gastroesophageal reflux in infants. Ital J Gastroenterol Hepatol, 1997; 29:237-42.
19. Wenzl TG et al. Effects of thickened feeding on gastroesophageal reflux in infants: a placebo-controlled crossover study using intraluminal impedance. Journal of Pediatrics, 2003;111(4):355-359
20. Laranjeira M et al. Adequado crescimento, ausência de ganho de peso excessivo e de alteração no padrão evacuatório em lactentes com refluxo gastroesofágico recebendo fórmula infantil espessada com goma jataí. Pediatr. Mod. 2014; 50(8): 339-342.
21. Simakachorn N et al. Randomized, double-blind clinical trial of a lactose-free and a lactose containing formula in dietary management of acute childhood diarrhea. J Med Assoc Thai. 2004 Jun;87(6):641- 9.
22. Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. Arq Asma Alerg Imunol. 2018;2(1):7-82.

23. Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.
24. Niggemann B, Binder C, Dupont C, Hadji S, Arvola T, Isolauri E. Prospective, controlled, multi-center study on the effect of an amino-acid-based formula in infants with cow's milk allergy/intolerance and atopic dermatitis. *Pediatr Allergy Immunol*. 2001;12(2):78-82.
25. Sampson et al (1992). Safety of an amino acid derived infant formula in children allergic to cow milk. *Pediatrics*, 90(3), 463 465.
26. Verwimp JJ, et al. Symptomatology and growth in infants with cow's milk protein intolerance using two different whey protein hydrolysate-based formulas in a Primary Health Care setting. *Eur J Clin Nutr*. 1995;49 Suppl 1: S39 S48.
27. Giampietro PG, et al. Hypoallergenicity of an extensively hydrolyzed whey formula. *Pediatr Allergy Immunol*. 2001;12(2):83 86.
28. Isolauri, E., et al (1995). Efficacy and safety of hydrolyzed cow milk and amino acid derived formulas in infants with cow milk allergy. *The Journal of pediatrics*, 127(4), 550 557.0.
29. Koletzko, Berthold, et al. "Scientific basis and practical application of nutritional care for preterm infants." *Nutritional Care of Preterm Infants*. Vol. 122. Karger Publishers, 2021.
30. Agostoni C et al. Enteral Nutrient Supply for Preterm Infants: Commentary from the European Society for Paediatric Gastroenterology, Hepatology, and Nutrition Committee on Nutrition. *Journal of Pediatric Gastroenterology and Nutrition*. 2010; 50 (1): 85-91

REFERÊNCIAS FÓRMULAS INFANTIS PARA PREMATUROS

1. Koletzko, Berthold, et al. "Scientific basis and practical application of nutritional care for preterm infants." *Nutritional Care of Preterm Infants*. Vol. 122. Karger Publishers, 2021.
2. Embleton, N. D., Moltu, S. J., Lapillonne et al. Enteral nutrition in preterm infants (2022): a position paper from the ESPGHAN committee on nutrition and invited experts. *Journal of pediatric gastroenterology and nutrition*.2023; 76(2), 248-268.
3. Aggett, P. J. Agostoni, C. Axelsson I. et al. 2006 Feeding preterm infants after hospital discharge: a commentary by the ESPGHAN Committee on Nutrition. *Journal of Pediatric Gastroenterology and Nutrition*, 42, 596-603.
4. Lapillonne A, Bronsky J, Campoy C, et al. Feeding the Late and Moderately Preterm Infant: A Position Paper of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr*. 2019;69(2):259-270.
5. Knol J et al. Increase of faecal bifidobacteria due to dietary oligosaccharides induces a reduction of clinically relevant pathogen germs in the faeces of formula-fed preterm infants. *Acta Paediatrica*, 2005; 94(Suppl 449): 31–33.
6. Bruzzese E et al. Early administration of GOS/FOS prevents intestinal and respiratory infections in infants. *J Pediatr Gastroenterol Nutr*. 2006; 42 (5): E95.
7. Comparativo realizado entre produtos da mesma categoria, em 04/2025, utilizando informações extraídas da tabela nutricional
8. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
9. Liu L et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. *J Lipid Res*. 2014;55(3):531-9
10. Bar-Yoseph F et al. Review of sn-2 palmitate oil implications for infant health. *Prostaglandins Leukot Essent Fatty Acids*. 2013;89(4):139-43
11. Roy CC et al. Correction of the malabsorption of the preterm infant with a medium-chain triglyceride formula. *J Pediatr*. 1975;86(3):446-50.
12. Koletzko, Berthold, et al. "Scientific basis and practical application of nutritional care for preterm infants." *Nutritional Care of Preterm Infants*. Vol. 122. Karger Publishers, 2021.
13. Embleton, N. D., Moltu, S. J., Lapillonne et al. Enteral nutrition in preterm infants (2022): a position paper from the ESPGHAN committee on nutrition and invited experts. *Journal of pediatric gastroenterology and nutrition*.2023; 76(2), 248-268.
14. Aggett, P. J. Agostoni, C. Axelsson I. et al. 2006 Feeding preterm infants after hospital discharge: a commentary by the ESPGHAN Committee on Nutrition. *Journal of Pediatric Gastroenterology and Nutrition*, 42, 596-603.

REFERÊNCIAS FÓRMULAS INFANTIS PARA LACTENTES

1. Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4^a ed. Rio de Janeiro, 2018.
2. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am.* 2013;60(1):49-74.
3. Koletzko B et al. Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *Am J Clin Nutr.* 2009; 89: 1836–45.
4. Dallas DC, Murray NM, Gan J. Proteolytic Systems in Milk: Perspectives on the Evolutionary Function within the Mammary Gland and the Infant. *J Mammary Gland Biol Neoplasia.* 2015 Dec;20(3-4):133-47.
5. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
6. Liu L et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. *J Lipid Res.* 2014;55(3):531-9
7. Bar-Yoseph F et al. Review of sn-2 palmitate oil implications for infant health. *Prostaglandins Leukot Essent Fatty Acids.* 2013;89(4):139-43
8. Salminen S, Szajewska H, Knol J. The Biotics Family in Early Life. Edited Wiley: vol. 4, 2019
9. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5
10. Bruzzese E et al. A formula containing galacto- and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study. *Clin Nutr.* 2009;28(2):156-61
11. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr.* 2013;98(2):561S-71S.
12. Scholtens PA et al. Stool characteristics of infants receiving short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides: a review. *World J Gastroenterol.* 2014;20(37):13446-52.
13. Vandenplas Y et al. A partly fermented infant formula with postbiotics including 3'-GL, specific oligosaccharides, 2'-FL, and milk fat supports adequate growth, is safe and well-tolerated in healthy term infants: a double-blind, randomised, controlled, multi-country trial. *Nutrients.* 2020; 12(11):3960
14. Overbeek S et al. Human milk oligosaccharide 2'-fucosyllactose more efficiently modulates immunogenicity during maturation of human dendritic cells in the presence of scGOS/IcFOS prebiotics. *JPGN.* 2019; 58:995
15. Comparativo realizado entre produtos da mesma categoria, em 04/2025, utilizando informações extraídas da tabela nutricional
16. Mun JG, Legette LL, Ikonte CJ, Mitmesser SH. Choline and DHA in Maternal and Infant Nutrition: Synergistic Implications in Brain and Eye Health. *Nutrients.* 2019 May 21;11(5):1125.

REFERÊNCIAS FÓRMULAS INFANTIS DE SEGUIMENTO PARA CRIANÇAS DE PRIMEIRA INFÂNCIA

1. Salminen S, Szajewska H, Knol J. The Biotics Family in Early Life. Edited Wiley: vol. 4, 2019
2. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5
3. Buzzese E et al. A formula containing galacto- and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study. *Clin Nutr.* 2009;28(2):156-61
4. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr.* 2013;98(2):561S-71S.
5. Scholtens PA et al. Stool characteristics of infants receiving short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides: a review. *World J Gastroenterol.* 2014;20(37):13446-52.
6. Luque et al. Early Programming by Protein Intake: The Effect of Protein on Adiposity Development and the Growth and Functionality of Vital Organs. *Nutrition and Metabolic Insights* 2015;8(S1) 49–56.
7. Koletzko B et al. Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *Am J Clin Nutr.* 2009; 89: 1836–45.
8. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
9. Mun JG, Legette LL, Ikonte CJ, Mitmesser SH. Choline and DHA in Maternal and Infant Nutrition: Synergistic Implications in Brain and Eye Health. *Nutrients.* 2019 May 21;11(5):1125.
10. Liu L et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. *J Lipid Res.* 2014;55(3):531-9
11. Bar-Yoseph F et al. Review of sn-2 palmitate oil implications for infant health. *Prostaglandins Leukot Essent Fatty Acids.* 2013;89(4):139-43

REFERÊNCIAS COMPOSTO LÁCTEO PARA CRIANÇAS A PARTIR DE 3 ANOS

1. Nogueira-de-Almeida, C. A., Falcão, M. C., Ribas Filho, D., Zorzo, R. A., Konstantyner, T., Ricci, R., Gioia, N., & Fisberg, M. (2020). Consensus of the Brazilian Association of Nutrology on Milky Feeding of Children Aged 1–5 Years Old. *International Journal of Nutrology*, 13(1), 2–16.
2. Luque et al. Early Programming by Protein Intake: The Effect of Protein on Adiposity Development and the Growth and Functionality of Vital Organs. *Nutrition and Metabolic Insights* 2015;8(S1) 49–56
3. Institute of Medicine (IOM). Dietary Reference Intakes: The Essential Guide to Nutrient Requirements, 2006.
4. Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4^a ed. Rio de Janeiro, 2018.
5. Nogueira-de-Almeida, C. A., Vilanova, K. C. M., Perini, T. M., Filho, D. R. (2024). Alimentação da Criança de Zero a Cinco Anos. Manual do Departamento de Nutrologia, ABRAN.
6. FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS. Fats and fatty acids in human nutrition: Report of an expert consultation. Rome: FAO, 2010.
7. Nogueira-de-Almeida CA, Ribas Filho D, Mello ED, Bertolucci PHF, Falcão MC. I Consenso da Associação Brasileira de Nutrologia sobre recomendações de DHA durante gestação, lactação e infância. In: Anais do XVIII Congresso Brasileiro de Nutrologia; 2014; São Paulo, BR. São Paulo; 2014. 1-13.”; 33.
8. Koletzko, B. (Berthold), editor. Pediatric Nutrition in Practice. 2nd, revised edition. (II. Series: World review of nutrition and dietetics; vol. 113); 2015
9. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306.
10. Cummings JH, Macfarlane GT. Gastrointestinal effects of prebiotics. *Br J Nutr.* 2002; 87 (Suppl 2):S145-S151.;
11. Veereman G. Pediatric applications of inulin and oligofructose. *J Nutr.* 2007; 137:2585S-9S.
12. Mello, Carolina Santos, Karina Vieira Barros, and Mauro Batista de Moraes, "Brazilian infant and preschool children feeding; literature review." *Jornal de pediatria* 92 (2016): 451-463.

REFERÊNCIAS DESAFIOS DO CRESCIMENTO

1. Kerzner B, Milano K, MacLean WC, et al. A Practical approach to classifying and managing feeding difficulties. 2015; 135(2):344-35
2. Nogueira CA, Mello E, Filho DR, Maximino P, Fisberg M. Consenso da Associação Brasileira de Nutrologia sobre o uso de suplementos alimentares para crianças com dificuldades alimentares. *Int J Nutrol* 2018;11(suppl S1):S4–S15
3. Sociedade Brasileira de Pediatria. Guia de orientações - Dificuldades alimentares. Departamento Científico de Nutrologia São Paulo: SBP, 2022.
4. Manual de Suporte Nutricional da Sociedade Brasileira de Pediatria - 2ed. - Rio de Janeiro: Departamento Científico de Suporte Nutricional da Sociedade Brasileira de Pediatria. – 2020.
5. Braegger C, Decsi T, Dias JA, Hartman C, Koláček S, Koletzko B et al. Practical Approach to Paediatric Enteral Nutrition: A Comment by the ESPGHAN Committee on Nutrition. *JPGN*; 2010;51(1):110-122
6. NICE Guideline. Faltering growth: recognition and management of faltering growth in children. 2017.
7. Mehta NM et al. Defining pediatric malnutrition: A paradigm shift toward etiology-related definitions. *J Parenter Enter Nutr.* 2013;37(4):460–81.
8. Codex Alimentarius FAO/OMS. Standard for infant formula and formulas for special medical purposes intended for infants. Codex Stan 72 – 1981. Revised 2007/2011.
9. Gahagan S. Failure to thrive: a consequence of undernutrition. *Pediatr Rev.* 2006 Jan;27(1):e1-11.
10. ASPEN. Nutrition Management of Term Infants with Growth Failure. 2022
11. Pencharz PB. Protein and energy requirements for “optimal” catch-up growth. *EJCN.*2010;64: S5-7
12. Clarke SE, Evans S, MacDonald A, Davies P, Booth IW. Randomized comparison of a nutrient-dense formula with an energy-supplemented formula for infants with faltering growth. *J Hum Nutr Diet.* 2007;20(4):329-339. 4
13. FAO/WHO/UNU. Protein and amino acid requirements in human nutrition: report of a joint FAO/WHO/UNU expert consultation. WHO technical report series no. 935, 2007.p.187
14. Falcão MC, Udsen N, Zamberlan P, Ceccon MER. *Rev Bras Nutr Clin* 2009; 24 (2): 125-30
15. Uauy R, Hoffman DR, Mena P, Llanos A, Birch EE. Term infant studies of DHA and ARA supplementation on neurodevelopment: results of randomized controlled trials. *J Pediatr.* 2003 ;143: S17-25.
16. SanGiovanni JP, Berkey CS, Dwyer JT, Colditz GA. Dietary essential fatty acids, long-chain polyunsaturated fatty acids, and visual resolution acuity in healthy full-term infants: a systematic review. *Early Hum Dev.* 2000; 57:165-88.
17. Knol J et al. Colon microflora in infants fed formula with galacto- and fructo-oligosaccharides: more like breast-fed infants. *J Pediatr Gastroenterol Nutr.* 2005;40(1):36-42.
18. Scholtens PA, Alles MS, Bindels JG, van der Linde EG, Tolboom JJ, Knol J. Bifidogenic effects of solid weaning foods with added prebiotic oligosaccharides: a randomised controlled clinical trial. *J Pediatr Gastroenterol Nutr.* 2006 May;42(5):553-9.
19. Food and Agriculture Organization of the United Nations. (2010). Fats and fatty acids in human nutrition. Report of an expert consultation. Food and Nutrition Paper 91.
20. Nogueira-de-Almeida CA, Ribas Filho D, Mello ED, Bertolucci PHF, Falcão MC. I Consenso da Associação Brasileira de Nutrologia sobre recomendações de DHA durante gestação, lactação e infância. In: Anais do XVIII Congresso Brasileiro de Nutrologia; 2014; São Paulo, BR. São Paulo; 2014. 1-13.”; 33.
21. Koletzko, B. (Berthold), editor. Pediatric Nutrition in Practice. 2nd, revised edition. (II. Series: World review of nutrition and dietetics; vol. 113); 2015
22. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306.
23. ILSI (International Life Sciences Institute) Brasil. Série: Funções plenamente reconhecidas de Nutrientes. 1 Ed. 2009-2014.
24. Devaera, Y., Syaharutsa, D. M., Jatmiko, H. K., & Sjarif, D. R. Comparing Compliance and Efficacy of Isocaloric Oral Nutritional Supplementation Using 1.5 kcal/mL or 1 kcal/mL Sip Feeds in Mildly to Moderately Malnourished Indonesian Children: A Randomized Controlled Trial. *Pediatric gastroenterology, hepatology & nutrition,* 21(4), 315–320.

25. Nogueira-de-Almeida, C.A.; Del Ciampo, L.A.; Martinez, E.Z.; Contini, A.A.; Nogueira-de-Almeida, M.E.; Ferraz, I.S.; Epifanio, M.; da Veiga Ued, F. Clinical Evolution of Preschool Picky Eater Children Receiving Oral Nutritional Supplementation during Six Months: A Prospective Controlled Clinical Trial. *Children* 2023, 10, 495.
26. Guideline: Sugars intake for adults and children. Geneva: World Health Organization; 2015.
27. Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4^a ed. Rio de Janeiro, 2018.
28. Nogueira-de-Almeida CA, Ribas Filho D, Mello ED, Bertolucci PHF, Falcão MC. I Consenso da Associação Brasileira de Nutrologia sobre recomendações de DHA durante gestação, lactação e infância. In: Anais do XVIII Congresso Brasileiro de Nutrologia; 2014; São Paulo, BR. São Paulo; 2014. 1-13.”; 33.
29. Scholtens PA et al. Stool characteristics of infants receiving short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides: a review. *World J Gastroenterol.* 2014;20(37):13446-52.
30. Knol J et al. Colon microflora in infants fed formula with galacto- and fructo-oligosaccharides: more like breast-fed infants. *J Pediatr Gastroenterol Nutr.* 2005;40():36-42.

REFERÊNCIAS DISTÚRBIOS GASTROINTESTINAIS FUNCIONAIS

1. Borrelli O et al. Use of a new thickened formula for treatment of symptomatic gastroesophageal reflux in infants. *Ital J Gastroenterol Hepatol*, 1997; 29:237-42.
2. Wenzl TG et al. Effects of thickened feeding on gastroesophageal reflux in infants: a placebo-controlled crossover study using intraluminal impedance. *Journal of Pediatrics*, 2003;111(4):355-359.
3. Rosen R, Vandenplas Y, Singendonk M, et al. Pediatric Gastroesophageal Reflux Clinical Practice Guidelines: Joint Recommendations of the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition and the European Society for Pediatric Gastroenterology, Hepatology, and Nutrition. *J Pediatr Gastroenterol Nutr*. 2018;66(3):516-554.
4. Salvatore, S. et al. Thickened infant formula: what to know. *Nutrition*, 2018.
5. Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation, and cow's milk allergy in formula-fed infants. *Acta Pædiatrica* 2015,
6. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am*.2013;60(1):49-74.Infante Pina D et al. Thickened infant formula, rheological study of the "in vitro" properties. *An Pediatr (Barc)*. 2010; 72(5):302-8.
7. Koletzko B et al. The roles of long-chain polyunsaturated fatty acids in pregnancy, lactation and infancy: review of current knowledge and consensus recommendations. *J Perinat Med*. 2008;36(1):5-14
8. Qawasmi A et al. Meta-analysis of LCPUFA supplementation of infant formula and visual acuity. *Pediatrics*. 2013;131(1):262-72.
9. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
10. Miles EA, Childs CE, Calder PC. Long-Chain Polyunsaturated Fatty Acids (LCPUFAs) and the Developing Immune System: A Narrative Review. *Nutrients*. 2021; 13: 247.
11. Lerner A, Shamir R. Nucleotides in infant nutrition: a must or an option. *Isr Med Assoc J*. 2000; 2(10):772-4.
12. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr*. 2002;34(3):291-5
13. Braegger C et al. Supplementation of infant formula with probiotics and/or prebiotics: a systematic review and comment by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr*. 2011;52:238–225
14. Boehm G et al. Prebiotics in infant formulas. *J Clin Gastroenterol*. 2004; 38: S76-9.
15. Vandenplas Y et al. Oligosaccharides in infant formula: more evidence to validate the role of prebiotics. *Br J Nutr*. 2015;113(9):1339-44.
16. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr*. 2013;98(2):561S-71S.
17. Savino F et al. Reduction of crying episodes owing to infantile colic: A randomized controlled study on the efficacy of a new infant formula. *Eur J Clin Nutr* 2006, 60:1304-10.
18. Savino F et al. "Minor" feeding problems during the first months of life: effect of a partially hydrolyzed milk formula containing fructo- and galacto-oligosaccharides. *Acta Paediatr Suppl* 2003, 91:86-90.
19. Schmelzle H et al. Randomized double-blind study of the nutritional efficacy and bifidogenicity of a new infant formula containing partially hydrolyzed protein, a high beta-palmitic acid level, and nondigestible oligosaccharides. *J Pediatr Gastroenterol Nutr* 2003, 36:343-51.
20. Bongers ME et al. The clinical effect of a new infant formula in term infants with constipation: a double-blind, randomized cross-over trial. *Nutr J* 2007;6:8.
21. Savino F et al. Advances in the management of digestive problems during the first months of life. *Acta Paediatr Suppl* 2005, 94:120.
22. Havlicekova Z et al. Beta-palmitate - a natural component of human milk in supplemental milk formulas. *Nutr J*. 2016;15(1):28.
23. Litmanovitz I et al. Reduced crying in term infants fed high beta-palmitate formula: a double-blind randomized clinical trial. *BMC Pediatr*. 2014;14:152.

24. Yao M et al. Effects of term infant formulas containing high sn-2 palmitate with and without oligofructose on stool composition, stool characteristics, and bifidogenicity. *J Pediatr Gastroenterol Nutr.* 2014 ;59(4):440-8.
25. Nowacki J et al. Stool fatty acid soaps, stool consistency and gastrointestinal tolerance in term infants fed infant formulas containing high sn-2 palmitate with or without oligofructose: a double-blind, randomized clinical trial. *Nutr J.* 2014;13:105.
26. Simakachorn N et al. Randomized, double-blind clinical trial of a lactose-free and a lactose containing formula in dietary management of acute childhood diarrhea. *J Med Assoc Thai.* 2004 Jun;87(6):641- 9.

REFERÊNCIAS ALERGIA À PROTEÍNA DO LEITE DE VACA (APLV)

1. Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. Arq Asma Alerg Imunol. 2018;2(1):7-82.
2. Koletzko S et al. Diagnostic approach and management of cow's-milk protein allergy in infants and children: ESPGHAN GI Committee practical guidelines. J Pediatr Gastroenterol Nutr. 2012;55 (2):221-9.
3. Muraro A et al. EAACI food allergy and anaphylaxis guidelines: diagnosis and management of food allergy. Allergy. 2014;69(8):1008-25.
4. Fiocchi A et al. World Allergy Organization (WAO) Diagnosis and Rationale for Action against Cow's Milk Allergy (DRACMA) Guidelines. Pediatr Allergy Immunol. 2010;21 Suppl 21:1-125.
5. Venter C et al. Diagnosis and management of non-IgE-mediated cow's milk allergy in infancy: a UK primary care practical guide. Clin Transl Allergy. 2013;3(1):23.
6. Venter C et al. Better recognition, diagnosis and management of non-IgE-mediated cow's milk allergy in infancy: iMAP-an international interpretation of the MAP (Milk Allergy in Primary Care) guideline. Clin Transl Allergy. 2017; 7:26.
7. Orsi M et al. Alergia a la proteína de la leche de vaca: Propuesta de Guía para el manejo de los niños con alergia a la proteína de la leche de vaca. Arch. argent. pediatr. 2009; 107 (5): 459-467.
8. Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.
9. Toca MC et al. Consenso sobre el diagnóstico y el tratamiento de la alergia a las proteínas de la leche de vaca de la Sociedad Latinoamericana de Gastroenterología, Hepatología y Nutrición. Rev Gastroenterol Méx. 2022;87:235---250.
10. Fox A, Brown T, Walsh J, et al. An update to the Milk Allergy in Primary Care guideline. Clin Transl Allergy. 2019;9:40. Published 2019 Aug 12.
11. Vandenplas Y, et al. An ESPGHAN position paper on the diagnosis, management and prevention of cow's milk allergy. J Pediatr Gastroenterol Nutr. 2023 Jul 26.
12. Morais MB, Spolidoro JV, Vieira MC, Cardoso AL, Clark O, Nishikawa A, Castro AP. Amino acid formula as a new strategy for diagnosing cow's milk allergy in infants: is it cost-effective? J Med Econ. 2016;19:1207-14.
13. Birch E. et al. The DIAMOND (DHA intake and measurement of neural development) study: a double-masked, randomized controlled clinical trial of the maturation of infant visual acuity as a function of the dietary level of docosahexaenoic acid' Am J Clin Nutr., vol. 91, no.4, pg. 848–859, 2010.
14. Qawasmi A et al. Meta-analysis of LCPUFA supplementation of infant formula and visual acuity. Pediatrics. 2013;131(1):262-72.
15. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. BRAS PEN J 2020; 35 (3): 294-306
16. Miles EA, Childs CE, Calder PC. Long-Chain Polyunsaturated Fatty Acids (LCPUFAs) and the Developing Immune System: A Narrative Review. Nutrients. 2021; 13: 247.
17. Marten B et al. Medium-chain triglycerides. International Dairy Journal. 2006;16: 1374-1382
18. Estudo observacional, prospectivo aberto e multicêntrico de vigilância pós-comercialização sobre aceitabilidade e tolerabilidade de uma fórmula infantil extensamente hidrolisada. Brasil, 2016: dados preliminares.
19. Epifanio M et al. Palatability and Sensory Perception of Infant Formulas for the Treatment of Cow's Milk Allergy According to Brazilian Mothers. Biomed J Sci & Tech Res 25(1)-2020. BJSTR. MS.ID.00416.
20. Eveleens, R D et al. Definitions, predictors, and outcomes of feeding intolerance in critically ill children: A systematic review. Clin Nutr, 2020, 39.3: 685-693.
21. Marino, LV et al. Feeding intolerance in children with critical illness. Clin. Nutr., 2020, 39.3: 609-611.
22. Lucia Diaferio L et al. May Failure to Thrive in Infants Be a Clinical Marker for the Early Diagnosis of Cow's Milk Allergy? Nutrients . 2020 Feb 13;12(2):466. doi: 10.3390/nu12020466.
23. Boaventura RM, Mendonça RB, Fonseca FA, Mallozi M, Souza FS, Sarni ROS. Nutritional status and food intake of children with cow's milk allergy. Allergol Immunopathol (Madr). 2019;47(6):544-550. doi:10.1016/j.aller.2019.03.003
24. Meyer R, Venter C, Fox AT, Shah N. Practical dietary management of protein energy malnutrition in young children with cow's milk protein allergy. Pediatr Allergy Immunol. 2012;23(4):307-314.
25. Nutrologia Pediátrica: Temas da Atualidade em Nutrologia Pediátrica - 2021. / Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia. São Paulo: SBP, 2021.

26. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am.* 2013;60(1):49-74
27. FERREIRA, CHT; BARRETO, BAP; TOFOLI, MHC; et al. Microbioma e pediatria na saúde e na doença. Barueri [SP]: Manole, 2024. 1.ed.
28. Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar/ Sociedade Brasileira de Pediatria. – 5.ed. rev. ampl.– São Paulo: SBP, 2024. 208 p.
29. Arslanoglu S, Moro GE, Boehm G, Wienz F, Stahl B, Bertino E. Early neutral prebiotic oligosaccharide supplementation reduces the incidence of some allergic manifestations in the first 5 years of life. *J Biol Regul Homeost Agents.* 2012;26(3 Suppl):49-59.
30. Arslanoglu S, Moro GE, Schmitt J, Tandoi L, Rizzardi S, Boehm G. Early Dietary intervention with a mixture of prebiotic oligosaccharides reduces the incidence of allergic manifestations and infections during the first two years of life. *J Nutr.* 2008;138(6):1091-5.
31. Boehm G. A mixture of prebiotics oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Arch Dis Child.* 2006;91(10):814-
32. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5
33. Koletzko B, Baker S, Cleghorn G, et al. Global standard for the composition of infant formula: recommendations of an ESPGHAN coordinated international expert group. *J Pediatr Gastroenterol Nutr.* 2005;41(5):584-99
34. SALMINEN, Seppo; SZAJEWSKA, Hania; KNOL, Jan. The Biotics Family in Early Life. 2. ed. Wiley, 2023.
35. Goh CY, et al. Combination of short-chain GOS and long-chain FOS 9:1 with 2FL positively impact the infant gut microbiota composition and metabolic activity in a stimulator of the human intestinal microbial ecosystem (SHIME). *J Pediatr Gastroenterol Nutr* 2019; 68 (S1): NP 114; 1159.
36. SBP. Guia de orientações - Dificuldades alimentares/ Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia. São Paulo: SBP, 2022. 66 f.
37. Medeiros LCS et al. Ingestão de nutrientes e estado nutricional de crianças em dieta isenta de leite de vaca e derivados. *J. Pediatr. (Rio J.)* vol.80 no.5 Porto Alegre, 2004
38. Meyer R et al. A practical approach to vitamin and mineral supplementation in food allergic children. *Clinical and Translational Allergy* (2015) 5:11
39. Meyer R, De Koker C, Dziubak R, Venter C, Dominguez-Ortega G, Cutts R et al. Malnutrition in children with food allergies in the UK. *J Hum Nutr Diet.* 2014;27(3):227-35
40. Veereman G. Pediatric applications of inulin and oligofructose. *J Nutr.* 2007; 137:2585S-9S.
41. HCP TRACKER, 2024

REFERÊNCIAS LINHA SOJA

1. Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. Arq Asma Alerg Imunol. 2018;2(1):7-82.
2. Bhatia J, Greer F. American Academy of Pediatrics Committee on Nutrition: Use of soy protein-based formulas in infant feeding. Pediatrics. 2008;121(5):1062-8
3. Agostoni C et al. Soy protein infant formulae and follow-on formulae: A commentary by the ESPGHAN Committee on Nutrition. J Ped Gast Nutr 2006; 42:352-61.
4. Vandenplas Y, Hegar B, Munasir Z, et al. The role of soy plant-based formula supplemented with dietary fiber to support children's growth and development: An expert opinion. Nutrition. 2021;90:111278.
5. Merritt RJ, Fleet SE, Fifi A, et al. North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition Position Paper: Plant-based Milks. J Pediatr Gastroenterol Nutr. 2020;71(2):276-281.
6. ILSI (International Life Sciences Institute) Brasil. Série: Funções plenamente reconhecidas de Nutrientes. 1 Ed. 2009-2014.
7. FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS. Fats and fatty acids in human nutrition: Report of an expert consultation. Rome: FAO, 2010. 7.
8. Nogueira-de-Almeida CA, Ribas Filho D, Mello ED, Bertolucci PHF, Falcão MC. I Consenso da Associação Brasileira de Nutrologia sobre recomendações de DHA durante gestação, lactação e infância. In: Anais do XVIII Congresso Brasileiro de Nutrologia; 2014; São Paulo, BR. São Paulo:; 2014. 1-13." ; 33.
9. Koletzko, B. (Berthold), editor. Pediatric Nutrition in Practice. 2nd, revised edition. (II. Series: World review of nutrition and dietetics ; vol. 113); 2015

REFERÊNCIAS ERROS INATOS DO METABOLISMO E NECESSIDADES ESPECÍFICAS

1. Sampaio LPB, Takakura C, Manreza, MLG. The use of a fórmula-based ketogenic diet in children with refractory epilepsy. Arq. NeuroPsiquiatr. 2017;75(4):234-237.
2. Weijenberg A, van Rijn M, Callenbach PMC, de Koning TJ, Brouwer OF. Ketogenic Diet in Refractory Childhood Epilepsy: Starting With a Liquid Formulation in an Outpatient Setting. Child Neurol Open. 2018; 5:2329048X18779497.
3. Thompson L, Fecske E, Salim M, Hall A. Use of the ketogenic diet in the neonatal intensive care unit-Safety and tolerability. Epilepsia. 2017;58(2):e36-e39.
4. Coppola G, Verrotti A, Ammendola E, Operto FF, Corte RD, Signoriello G et al. Ketogenic diet for the treatment of catastrophic epileptic encephalopathies in childhood. Eur J Paediatr Neurol. 2010; 14(3):229–234.
5. Kossoff EH, Dorward JL, Turner Z, Pyzik PL. Prospective study of the modified atkins diet in combination with a ketogenic liquid supplement during the initial month. J Child Neurol. 2011;26(2):147-51.
6. Ashrafi MR Hosseini SA, Zamani GR, Mohammadi M, Tavassoli A, Badv RS et al. The efficacy of the ketogenic diet in infants and Young children with refractory epilepsies using a formula-based powder. Acta Neurol Belg. 2017;117(1):175-182.
7. Uldall P et al. Ketogenic diet for infants is possible with Ketocal. Epilepsia. 2006;47(S3):176.
8. Rubenstein JE. Use of the ketogenic diet in neonates and infants. Epilepsia. 2008;49(Suppl 8):30.
9. Nizamuddin J, Turner Z, Rubenstein JE, Pyzik PL, Kossoff EH. Management and risk factors for dyslipidemia with the ketogenic diet. J Child Neurol. 2008;23(7):758-61].
10. Kossoff EH, Zupec- Kania BA, Amark PE, Ballaban-Gil KR, Christina Bergqvist AG, Blackford R et al. Optimal Clinical Management of Children receiving the ketogenic diet: recommendations of the International ketogenic diet study group. Epilepsia. 2009;50(2):304- 317.
11. Sampaio LPB. ABC da dieta cetogênica para epilepsia refratária. Rio de Janeiro: Editora DOC Content, 2018. 1a edição,220 p.
12. Zamberlan P et al. Nutrição enteral em pediatria. Pediatria Moderna. 2002; 38(4):105-124.
13. Van der Louw E et al. Ketogenic diet guidelines for infants with refractory epilepsy. Eur J Paediatr Neurol. 2016 Nov;20(6):798-809.