

## REFERÊNCIAS BIBLIOGRÁFICAS

### Calendário Institucional 2024

1. Supinski GS, Schroder EA, Callahan LA. Mitochondria and Critical Illness. *Chest.* 2020 Feb;157(2):310-322.
2. Galley HF. Oxidative stress and mitochondrial dysfunction in sepsis. *Br J Anaesth.* 2011 Jul;107(1):57-64.
3. Amengual J. Bioactive Properties of Carotenoids in Human Health. *Nutrients.* 2019 Oct 6;11(10):2388
4. Quasim T, McMillan DC, Talwar D, Sattar N, O'Reilly DS, Kinsella J. Lower concentrations of carotenoids in the critically ill patient are related to a systemic inflammatory response and increased lipid peroxidation. *Clin Nutr.* 2003 Oct;22(5):459-62.
5. Wanten G, Calder P. Immune modulation by parenteral lipid emulsions. *Am J Clin Nutr* 2007;85:1171–84.
6. Vaisman et al. Enteral feeding enriched with carotenoids normalizes the carotenoid status and reduces oxidative stress in long-term enterally fed patients. *Clinical Nutrition* (2006) 25, 897–905
7. Liu J, Klebach M, Visser M, Hofman Z. Amino Acid Availability of a Dairy and Vegetable Protein Blend Compared to Single Casein, Whey, Soy, and Pea Proteins: A Double-Blind, Cross-Over Trial. *Nutrients.* 2019 Nov 1;11(11):2613.
8. Braak CCvd, Klebach M, Abrahamse E, et al. A novel protein mixture containing vegetable proteins renders enteral nutrition products non-coagulating after in vitro gastric digestion. *Clinical nutrition (Edinburgh, Scotland)* 2013; 32(5): 765-71. 9. Liu J, et al. Specific protein mixture reduces coagulation: An in vitro stomach model study mimicking a gastric condition in critically ill patients. *ESPEN 2016* 2016; MON-P182
10. Goelen N, Janssen P, Ripken D, van Horssen P, Byloos K, Ghysels S, Putzeys G, Hofman Z, Vandecaveye V, Tack J. Effect of protein composition of enteral formula on gastric content volume during continuous feeding: A randomized controlled cross-over study in healthy adults. *Clin Nutr.* 2021 May;40(5):2663-2672.
11. Weijs PJ, et al. Optimal protein and energy nutrition decreases mortality in mechanically ventilated, critically ill patients: A prospective observational cohort study. *JPEN,* 2012;36(1):60-8.
12. Allingstrup MJ, et al. Provision of protein and energy in relation to measured requirements in intensive care patients. *Clin Nutr,* 2012;31(4):462-8.
13. Roy et al. New lipids in enteral feeding. *Curr Opin Clin Nutr Metab Care* (2004) 7:117– 122.
14. Waitzberg, DL. Nutrição oral, enteral e parenteral na prática clínica/ Dan L. Waitzberg.- 4edição-São Paulo: Editora Atheneu,2009
15. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306. 28
16. Barazzoni R, Deutz NEP, Biolo G, et al. Carbohydrates and insulin resistance in clinical nutrition: Recommendations from the ESPEN expert group. *Clin Nutr.* 2017;36(2):355-363.
17. Yagmurdur H, Leblebici F. Enteral nutrition preference in critical care: fibre-enriched or fibre-free? *Asia Pac J Clin Nutr.* 2016 Dec;25(4):740-746.
18. Jakobsen LH, Wirth R, Smoliner C, Klebach M, Hofman Z, Kondrup J. Gastrointestinal tolerance and plasma status of carotenoids, EPA and DHA with a fiber-enriched tube feed in hospitalized patients initiated on tube nutrition: Randomized controlled trial. *Clin Nutr.* 2017 Apr;36(2):380-388.
19. Daly A, Johnson T, MacDonald A. Is fibre supplementation in paediatric sip feeds beneficial? *J Hum Nutr Diet.* 2004 Aug;17(4):365-70.
20. Hofman Z., et al. Tolerance and efficacy of a multi-fibre enriched tube-feed in paediatric burn patients. *Clinical Nutrition.* 2001. 20 (3), abstract 217. 21. Schneider SM., et al. The effect of a polymeric enteral formula supplemented with a mixture of six fibres on normal human bowel function and colonic motility. *Clinical Nutrition,* 2006. 25(1): 82-90.
22. Guimber D., et al. Effect of multifibre mixture with prebiotic components on bifidobacterial and stool pH in tube-fed children. *British Journal of Nutrition,* 2010. 104(10), 1514-1522.
23. Karakan T., et al. Comparison of early enteral nutrition in severe acute pancreatitis with prebiotic fiber supplementation versus standard enteral solution: a prospective randomized double-blind study. *World J Gastr,* 2007. 13(19): 2733-2737

- 24.** Burgos R et al. ESPEN guideline clinical nutrition in neurology. *Clinical Nutrition* 37 (2018) 354e396
- 25.** SILVA, R R et al. Efeito da utilizacao de gomas na viscosidade e nas caracteristicas sensoriais de shake a base de farinha de banana verde. *Braz. J. Food Technol.* [online]. 2018, vol.21 [citado 2019-09-30], e2016052.
- 26.** Bauer, J. et al. Evidence-Based Recommendations for Optimal Dietary Protein Intake in Older People: A Position PaperFrom the PROT-AGE Study Group. *J Am Med Dir Assoc.* 2013 Aug;14(8):542-59 27.
- Bauer JM et al. Effects of a Vitamin D and Leucine-Enriched Whey Protein Nutritional Supplement on Measures of Sarcopenia in Older Adults, the PROVIDE Study: A Randomized, Double-Blind, Placebo-Controlled Trial. *J Am Med Dir Assoc.* 2015 Sep 1;16(9):740- 7.
- 28.** Paddon-Jones D. et al. *Curr Opin Clin Nutr Metab Care.* 2009 Jan;12(1):86-9
- 29.** Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyere O, Cederholm T, et al., Writing Group for the European Working Group on Sarcopenia in Older People 2 (EWGSOP2), and the Extended Group for EWGSOP2. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing.*
- 30.** Morley, J. et al. Frailty Consensus: a call to action. *J Am Med Dir Assoc* 2013, June; 14(6): 392–397
- 31.** Lombard K, et al. Compliance of energy-dense, small volume oral nutritional supplements in the daily clinical practice on a geriatric Ward—an observational study. *J Nutr Health Aging,* 2014;18(7):649-653.
- 32.** Hubbard GP, et al. A systematic review of compliance to oral nutritional supplements. *Clin nutri,* 2012;31(3):293-312
- 33.** Volkert D et al. ESPEN Guidelines on Enteral Nutrition: Geriatrics. *Clin Nutr.* 2006 Apr;25(2):330-60.
- 34.** Kreymann KG, Berger MM et al. ESPEN Guidelines on Enteral Nutrition: Intensive care. *Clin Nutr.* 2006 Apr;25(2):210-23
- 35.** Matos LBN et al. Campanha Diga Nao a Lesao por Pressao. *BRASPEN J* 2020; 35 (Supl 1):2-32.
- 36.** European Pressure Ulcer Advisory Panel - EPUAP - Prevencao e tratamento de ulcera por pressao / lesoes: pratica clinica Diretriz. A Diretriz Internacional. Emily Haesler (Ed.). EPUAP / NPIAP / PPPIA: 2019
- 37.** Diretriz Braspen de Terapia Nutricional no Envelhecimento. *BRASPEN* 2019;34(Supl3):2-58.
- 38.** Zambelli CMSF, et al. Diretriz BRASPEN de Terapia Nutricional no Paciente com Doenca Renal. *BRASPEN J* 2021; 36 (2o Supl 2): 2-22.
- 39.** Fiacchadori E, et al. ESPEN guideline on clinical nutrition in hospitalized patients with acute or chronic kidney disease. *Clinical Nutrition* 40 (2021) 1644e1668.
- 40.** Ikizler TA, Cuppari L. The 2020 Updated KDOQI Clinical Practice Guidelines for Nutrition in Chronic Kidney Disease. *Blood Purif* 2021;50:667–671
- 41.** Sampaio LPB, Takakura C, Manreza, MLG. The use of a formula-based ketogenic diet in children with refractory epilepsy. *Arq. NeuroPsiquiatr.* 2017;75(4):234-237.
- 42.** Weijenberg A, van Rijn M, Callenbach PMC, de Koning TJ, Brouwer OF. Ketogenic Diet in Refractory Childhood Epilepsy: Starting With a Liquid Formulation in an Outpatient Setting. *Child Neurol Open.* 2018;5:2329048X18779497.
- 43.** Thompson L, Fecske E, Salim M, Hall A. Use of the ketogenic diet in the neonatal intensive care unit-Safety and tolerability. *Epilepsia.* 2017;58(2):e36-e39.
- 44.** Coppola G, Verrotti A, Ammendola E, Operto FF, Corte RD, Signoriello G et al. Ketogenic diet for the treatment of catastrophic epileptic encephalopathies in childhood. *Eur J Paediatr Neurol.* 2010; 14(3):229–234.
- 45.** Kossoff EH, Dorward JL, Turner Z, Pyzik PL. Prospective study of the modified atkins diet in combination with a ketogenic liquid supplement during the initial month. *J Child Neurol.* 2011;26(2):147-51.
- 46.** Ashrafi MR, Hosseini SA, Zamani GR, Mohammadi M, Tavassoli A, Badv RS et al. The efficacy of the ketogenic diet in infants and young children with refractory epilepsies using a formula-based powder. *Acta Neurol Belg.* 2017;117(1):175-182.
- 47.** Uldall P et al. Ketogenic diet for infants is possible with Ketocal. *Epilepsia.* 2006;47(S3):176.
- 48.** Rubenstein JE. Use of the ketogenic diet in neonates and infants. *Epilepsia.* 2008;49(Suppl 8):30.

- 49.** Nizamuddin J, Turner Z, Rubenstein JE, Pyzik PL, Kossoff EH. Management and risk factors for dyslipidemia with the ketogenic diet. *J Child Neurol*. 2008;23(7):758-61].
- 50.** Newton CR, Epilepsy in poor regions of the world *The Lancet*, Volume 380, Issue 9848;1193-1201, 2012.
- 51.** Kossoff EH et al. Optimal clinical management of children receiving dietary therapies for epilepsy: Updated recommendations of the International Ketogenic Diet Study Group. *Epilepsia Open*. 2018 Jun; 3(2): 175–192
- 52.** Kossoff E et al. When do seizures usually improve with the ketogenic diet? *Epilepsia*. 2008;49:329-333