

REFERÊNCIAS BIBLIOGRÁFICAS

Referências – Lâmina Ganepão 2024

1. Liu J, et al. Amino Acid Availability of a Dairy and Vegetable Protein Blend Compared to Single Casein, Whey, Soy, and Pea Proteins: A Double-Blind, Cross-Over Trial. *Nutrients.* 2019 Nov 1;11(11):2613
2. Goelen N, et al. Effect of protein composition of enteral formula on gastric content volume during continuous feeding: A randomized controlled cross-over study in healthy adults. *Clin Nutr.* 2021 May;40(5):2663- 2672.
3. Braak CC, et al. A novel protein mixture containing vegetable proteins renders enteral nutrition products non-coagulating after in vitro gastric digestion. *Clinical nutrition (Edinburgh, Scotland)* 2013; 32(5):765-71.
4. Liu J, et al. Specific protein mixture reduces coagulation: An in vitro stomach model study mimicking a gastric condition in critically ill patients. *ESPEN 2016* 2016; MON-P182
5. Marlett, JA. et al. Position of the American Dietetic Association: health implications of dietary fiber. *J. Am Dietetic Assoc.* 102 (7), 2002, 993-1000.
6. Ilsi Brasil. Funções plenamente reconhecidas. Disponível em: <http://ilsi.org/publication/funcoes-plenamente-reconhecidas/>. Jan 2014.
7. Bernaud FSR; Rodrigues, TC. Fibra Alimentar – Ingestão adequada e efeitos sobre a saúde do metabolismo. *Arq Bras Endocrinol Metab.* 2013;57/6
8. Yagmurdur H, Leblebici F. Enteral nutrition preference in critical care: fibre-enriched or fibre-free? *Asia Pac J Clin Nutr.* 2016 Dec;25(4):740-746.
9. Jakobsen LH, Wirth R, Smoliner C, Klebach M, Hofman Z, Kondrup J. Gastrointestinal tolerance and plasma status of carotenoids, EPA and DHA with a fiber-enriched tube feed in hospitalized patients initiated on tube nutrition: Randomized controlled trial. *Clin Nutr.* 2017 Apr;36(2):380-388
10. Daly A, Johnson T, MacDonald A. Is fibre supplementation in paediatric sip feeds beneficial? *J Hum Nutr Diet.* 2004 Aug;17(4):365-70. doi: 10.1111/j.1365-277X.2004.00535.x. PMID: 15250846.
11. Roy et al. New lipids in enteral feeding. *Curr Opin Clin Nutr Metab Care* (2004) 7:117–122.
12. Waitzberg, DL. Nutrição oral, enteral e parenteral na prática clínica/ Dan L. Waitzberg.- 4edição-São Paulo: Editora Atheneu,2009
13. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306.
14. Supinski GS, Schroder EA, Callahan LA. Mitochondria and Critical Illness. *Chest.* 2020 Feb;157(2):310-322.
15. Quasim T, McMillan DC, Talwar D, Sattar N, O'Reilly DS, Kinsella J. Lower concentrations of carotenoids in the critically ill patient are related to a systemic inflammatory response and increased lipid peroxidation. *Clin Nutr.* 2003 Oct;22(5):459-62.
16. Galley HF. Oxidative stress and mitochondrial dysfunction in sepsis. *Br J Anaesth.* 2011 Jul;107(1):57-64.
17. Amengual J. Bioactive Properties of Carotenoids in Human Health. *Nutrients.* 2019 Oct 6;11(10):2388
18. Bauer, J. et al. Evidence-Based Recommendations for Optimal Dietary Protein Intake in Older People: A Position Paper From the PROT-AGE Study Group. *J Am Med Dir Assoc.* 2013 Aug;14(8):542-59

19. Bauer JM et al. Effects of a Vitamin D and Leucine-Enriched Whey Protein Nutritional Supplement on Measures of Sarcopenia in Older Adults, the PROVIDE Study: A Randomized, Double-Blind, Placebo-Controlled Trial. *J Am Med Dir Assoc.* 2015 Sep 1;16(9):740- 7.
20. Paddon-Jones D. et al. *Curr Opin Clin Nutr Metab Care.* 2009 Jan;12(1):86-90. 14. Cruz-Jentoft AJ. et al. *Age Ageing.* 2019 Jan 1;48(1):16-31.
21. Morley, J. et al. Frailty Consensus: a call to action. *J Am Med Dir Assoc* 2013, June; 14(6): 392–397.
22. Vaillant MF, et al. Guidelines on Standard and Therapeutic Diets for Adults in Hospitals by the French Association of Nutritionist Dieticians (AFDN) and the French Speaking Society of Clinical Nutrition and Metabolism (SFNCM). *Nutrients.* 2021 Jul 15;13(7):2434.
23. Kandiah et.al.,2006. Visual plate waste in hospitalized patients: length of stay and diet order. *J Am Diet Assoc*, v. 106, n. 10, p. 1663 – 1666.
24. Malafarina et.al.,2013.The anorexia of ageing: physiopathology, prevalence, associated comorbidity and mortality. A systematic review. *Maturitas.* Apr;74(4):293-302.
25. Norman K, Pichard C, Lochs H, Pirllich M. Prognostic impact of disease-related malnutrition. *Clin Nutr.* 2008 Feb;27(1):5-15.
26. Waitzberg DL, Caiaffa WT, Correia MI. Hospital malnutrition: the Brazilian national survey (IBRANUTRI): a study of 4000 patients. *Nutrition.* 2001 Jul-Aug;17(7-8):573-80.
27. Correia MITD, Perman MI, Waitzberg DL. Hospital malnutrition in Latin America: A systematic review. *Clin Nutr.* 2017 Aug;36(4):958-967.
28. Matos LBN et al. Campanha Diga Não à Lesão por Pressão. *BRASPEN J* 2020; 35 (Supl 1):2-32.
29. European Pressure Ulcer Advisory Panel - EPUAP - Prevenção e tratamento de úlceras por pressão / lesões: prática clínica Diretriz. A Diretriz Internacional. Emily Haesler (Ed.). EPUAP / NPIAP / PPPIA: 2019.
30. Mendes D, et al. A importância da nutrição no processo de Cicatrização de Feridas. *Revista Científica Univiçosa.* 2017; 9(1).
31. Meyers NA. et al. Nutrient support of the healing wound. *New Horiz.* 1994; 2(2):202-214.
32. Scholl D et. al. Nutrient recommendations for wound healing. *Journal of Intravenous Nursing.* 2001; 24 (2): 124-132.
33. Diretrizes da Sociedade Brasileira de Diabetes 2019-2020.
34. Vaisman N, et al. Tube feeding with a diabetes specific feed for 12 weeks improves glycaemic control in type 2 diabetes patients. *Clin Nutr.* 2009 Oct;28(5):549-55.
35. BRASPEN (2018) - Diretrizes Brasileiras de Nutrição Parenteral e Enteral no Paciente Grave - *BRASPEN J* 2018; 33 (Supl 1):2-36.
36. Brown B., et al. Enteral Nutrition Formula Selection: Current Evidence and Implications for Practice Nutrition in Clinical Practice, 2015; 30(1): 72–85.
37. Avenell A, Handoll HH. Nutritional supplementation for hip fracture aftercare in the elderly. *Cochrane Database Syst Rev* 2005; Issue 2. D001880. pub3.
38. Stratton RJ, Marinos E. A review of reviews: A new look at the evidence for oral nutritional supplements in clinical practice. *Clinical Nutrition Supplements,* 2007;2: 5–23.
39. Hetherington, M.M. Older adults and patients in need of nutritional support: Review of current treatment options and factors influencing nutritional intake. *Clinical Nutrition* 29 (2010) 160–169

- 40.** SILVA, R R et al. Efeito da utilização de gomas na viscosidade e nas características sensoriais de shake à base de farinha de banana verde. *Braz. J. Food Technol.* [online]. 2018, vol.21 [citado 2019-09-30], e2016052
- 41.** Oluseyi A, Enajite O. Malnutrition in Pre-Dialysis Chronic Kidney Disease Patients in a Teaching Hospital in Southern Nigeria. *Afr Health Sci.* 2016 Mar;16(1):234-41.
- 42.** Carrero JJ, Thomas F, Nagy K, Arogundade F, Avesani CM, Chan M, et al. Global Prevalence of Protein-Energy Wasting in Kidney Disease: A Meta-Analysis of Contemporary Observational Studies From the International Society of Renal Nutrition and Metabolism. *J Ren Nutr.* 2018 Nov;28(6):380-92.
- 43.** Obi Y, Qader H, Kovesdy CP, Kalantar-Zadeh K. Latest Consensus and Update on Protein-Energy Wasting in Chronic Kidney Disease. *Curr Opin Clin Nutr Metab Care.* 2015 May;18(3):254-62.
- 44.** Wu HL, Sung JM, Kao MD, Wang MC, Tseng CC, Chen ST. Nonprotein calorie supplement improves adherence to low-protein diet and exerts beneficial responses on renal function in chronic kidney disease. *J Ren Nutr.* 2013 Jul;23(4):271-6.