

Referências Multiplicando Experiências Caso Clínico

1. Lacono G, et al; Pediatric Study Group on Gastrointestinal Symptoms in Infancy. Gastrointestinal symptoms in infancy: a population-based prospective study. *Dig Liver Dis.* 2005 Jun;37(6):432-8;
2. Vandenplas Y, et al. Prevalence and Health Outcomes of Functional Gastrointestinal Symptoms in Infants From Birth to 12 Months of Age. *J Pediatr Gastroenterol Nutr.* 2015 Nov;61(5):531-7. doi: 10.1097/MPG.0000000000000949. Erratum in: *J Pediatr Gastroenterol Nutr.* 2016 Mar;62(3):516.;
3. Bellaiche M, et al. Multiple functional gastrointestinal disorders are frequent in formula-fed infants and decrease their quality of life. *Acta Paediatr.* 2018 Jul;107(7):1276-1282.;
4. Zeevenhooven J, Koppen IJ, Benninga MA. The New Rome IV Criteria for Functional Gastrointestinal Disorders in Infants and Toddlers. *Pediatr Gastroenterol Hepatol Nutr.* 2017 Mar;20(1):1-13.;
5. de Moraes MB, Toporovski MS, Tofoli MHC, Barros KV, Silva LR, Ferreira CHT. Prevalence of Functional Gastrointestinal Disorders in Brazilian Infants Seen in Private Pediatric Practices and Their Associated Factors. *J Pediatr Gastroenterol Nutr.* 2022 Jul 1;75(1):17-23.;
6. Vandenplas Y, Hauser B, Salvatore S. Functional Gastrointestinal Disorders in Infancy: Impact on the Health of the Infant and Family. *Pediatr Gastroenterol Hepatol Nutr.* 2019 May;22(3):207-216. doi: 10.5223/pghn.2019.22.3.207.;
7. Rosen R, et al. Pediatric Gastroesophageal Reflux Clinical Practice Guidelines: Joint Recommendations of the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition and the European Society for Pediatric Gastroenterology, Hepatology, and Nutrition. *J Pediatr Gastroenterol Nutr.* 2018 Mar;66(3):516-554. doi: 10.1097/MPG.0000000000001889.;
8. Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation and cow's milk allergy in formula-fed infants. *Acta Paediatrica* 2015, ISSN 0803-5253)
9. Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. *Arq Asma Alerg Imunol.* 2018;2(1):7-82.;
10. Vandenplas Y, et al. An ESPGHAN position paper on the diagnosis, management and prevention of cow's milk allergy. *J Pediatr Gastroenterol Nutr.* 2023 Jul 26.;
11. Atualização em Alergia Alimentar 2025: posicionamento conjunto da ASBAI e SBP – de Oliveira LCL, et al. *Arq Asma Alerg Imunol – Vol. 9, Nº 1, 2025;*
12. Savino F et al. Reduction of crying episodes owing to infantile colic: A randomized controlled study on the efficacy of a new infant formula. *Eur J Clin Nutr* 2006, 60:1304-10.;
13. Savino F et al. "Minor" feeding problems during the first months of life: effect of a partially hydrolyzed milk formula containing fructo- and galacto-oligosaccharides. *Acta Paediatr Suppl* 2003, 91:86-90.;
14. Schmelzle H et al. Randomized double-blind study of the nutritional efficacy and bifidogenicity of a new infant formula containing partially hydrolyzed protein, a high beta-palmitic acid level, and nondigestible oligosaccharides. *J Pediatr Gastroenterol Nutr* 2003, 36:343-51.;
15. Bongers ME et al. The clinical effect of a new infant formula in term

infants with constipation: a double-blind, randomized cross-over trial. *Nutr J* 2007;6:8.; 15. Savino F et al. Advances in the management of digestive problems during the first months of life. *Acta Paediatr Suppl* 2005, 94:120.; 16. Borrelli O et al. Use of a new thickened formula for treatment of symptomatic gastroesophageal reflux in infants. *Ital J Gastroenterol Hepatol*, 1997; 29:237-42.; 17. Wenzl TG et al. Effects of thickened feeding on gastroesophageal reflux in infants: a placebo-controlled crossover study using intraluminal impedance. *Journal of Pediatrics*, 2003;111(4):355-359.; 18. Salvatore, S. et al. Thickened infant formula: what to know. *Nutrition*, 2018.; 19. Infante Pina D et al. Thickened infant formula, rheological study of the “in vitro” properties. *An Pediatr (Barc)*. 2010; 72(5):302-8.; 20. Horvath A, Dziechciarz P, Szajewska H. The Effect of Thickened-Feed Interventions on Gastroesophageal Reflux in Infants: Systematic Review and Meta-analysis of Randomized, Controlled Trials. *Pediatrics* 2008;122:e1268–e1277.; 21. Simakachorn N et al. Randomized, double-blind clinical trial of a lactose-free and a lactose containing formula in dietary management of acute childhood diarrhea. *J Med Assoc Thai*. 2004 Jun;87(6):641- 9.; 22. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr*. 2002;34(3):291-5; 23. Braegger C et al. Supplementation of infant formula with probiotics and/or prebiotics: a systematic review and comment by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr*. 2011;52:238–225; 24. Boehm G et al. Prebiotics in infant formulas. *J Clin Gastroenterol*. 2004; 38: S76-9; 25. Vandenplas Y et al. Oligosaccharides in infant formula: more evidence to validate the role of prebiotics. *Br J Nutr*. 2015;113(9):1339-44; 26. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr*. 2013;98(2):561S-71S; 27. Havlicekova Z et al. Beta-palmitate - a natural component of human milk in supplemental milk formulas. *Nutr J*. 2016;15(1):28; 28. Litmanovitz I et al. Reduced crying in term infants fed high beta-palmitate formula: a double-blind randomized clinical trial. *BMC Pediatr*. 2014;14:152; 29. Yao M et al. Effects of term infant formulas containing high sn-2 palmitate with and without oligofructose on stool composition, stool characteristics, and bifidogenicity. *J Pediatr Gastroenterol Nutr*. 2014 ;59(4):440-8; 30. Nowacki J et al. Stool fatty acid soaps, stool consistency and gastrointestinal tolerance in term infants fed infant formulas containing high sn-2 palmitate with or without oligofructose: a double-blind, randomized clinical trial. *Nutr J*. 2014;13:105; 31. Sampaio LPB, Takakura C, Manreza, MLG. The use of a formula-based ketogenic diet in children with refractory epilepsy. *Arq. NeuroPsiquiatri*. 2017;75(4):234-237; 32. Weijenberg A, van Rijn M, Callenbach PMC, de Koning TJ, Brouwer OF. Ketogenic Diet in Refractory Childhood Epilepsy: Starting With a Liquid Formulation in an Outpatient Setting. *Child Neurol Open*. 2018;5:2329048X18779497; 33. Uldall P et al. Ketogenic diet for infants is possible

with Ketocal. *Epilepsia*. 2006;47(S3):176; 34. Nizamuddin J, Turner Z, Rubenstein JE, Pyzik PL, Kossoff EH. Management and risk factors for dyslipidemia with the ketogenic diet. *J Child Neurol*. 2008;23(7):758-61]; 35. Kossoff EH, Zupec-Kania BA, Amark PE, Ballaban-Gil KR, Christina Bergqvist AG, Blackford R et al. Optimal Clinical Management of Children receiving the ketogenic diet: recommendations of the International ketogenic diet study group. *Epilepsia*. 2009;50(2):304-317.; 36. Zamberlan P et al. Nutrição enteral em pediatria. *Pediatria Moderna*. 2002; 38(4):105-124.