

ONE PED DANONE

- 1.Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4^a ed. Rio de Janeiro, 2018.
- 2.Zubler et al. Evidence-informed milestones for developmental surveillance tools Pediatrics. 2022;149 (3)
- 3.Nogueira-de-Almeida, C. A., Falcão, M. C., Ribas Filho, D., Zorzo, R. A., Konstantyner, T., Ricci, R., Gioia, N., & Fisberg, M. (2022). Consensus of the Brazilian Association of Nutrology on Milky Feeding of Children Aged 1–5 Years Old. International Journal of Nutrology, 13(1), 2–16.
- 4.Manual de Suporte Nutricional da Sociedade Brasileira de Pediatria - 2ed. - Rio de Janeiro: Departamento Científico de Suporte Nutricional da
- 5.Sociedade Brasileira de Pediatria. – 2020. SBV. Pesquisa do IBOPE aponta crescimento histórico no número de vegetarianos no Brasil, Disponível em: Pesquisa do IBOPE aponta crescimento histórico no número de vegetarianos no Brasil - SBV. Acesso em 01.03.24.
- 6.Savilahti et al. 2012. 29.
- 7.Merritt, R. et al. North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition Position Paper: Plantbased Milks. JPGN Volume 71, Number 2, August 2020.
- 8.Sociedade Brasileira de Pediatria. Guia de orientações - Dificuldades alimentares. Departamento Científico de Nutrologia São Paulo: SBP, 2022.
- 9.Codex Alimentarius FAO/OMS. Standard for infant formula and formulas for special medical purposes intended for infants. Codex Stan 72 – 1981. Revised 2007/2011.
- 10.Gahagan S. Failure to thrive: a consequence of undernutrition. Pediatr Rev. 2006 Jan;27(1):e1-11.
- 11.Benninga MA, Faure C, Hyman PE, et al. Childhood functional gastrointestinal disorders:neonate/toddler. Gastroenterology 2016;150:1443-55.
- 12.Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation and cow's milk allergy in formula-fed infants. Acta Pædiatrica 2015, ISSN 0803-5253

- 13.Savino F et al. Reduction of crying episodes owing to infantile colic: A randomized controlled study on the efficacy of a new infant formula. Eur J Clin Nutr 2006, 60:1304-10.
- 14.Savino F et al. "Minor" feeding problems during the first months of life: effect of a partially hydrolyzed milk formula containing fructo- and galacto-oligosaccharides. Acta Paediatr Suppl 2003, 91:86-90.
- 15.Schmelzle H et al. Randomized double-blind study of the nutritional efficacy and bifidogenicity of a new infant formula containing partially hydrolyzed protein, a high beta-palmitic acid level, and nondigestible oligosaccharides. J Pediatr Gastroenterol Nutr 2003, 36:343-51.
- 16.Bongers ME et al. The clinical effect of a new infant formula in term infants with constipation: a double-blind, randomized cross-over trial. Nutr J 2007;6:8.
- 17.Savino F et al. Advances in the management of digestive problems during the first months of life. Acta Paediatr Suppl 2005, 94:120.
- 18.Borrelli O et al. Use of a new thickened formula for treatment of symptomatic gastroesophageal reflux in infants. Ital J Gastroenterol Hepatol, 1997; 29:237-42.
- 19.Wenzl TG et al. Effects of thickened feeding on gastroesophageal reflux in infants: a placebo-controlled crossover study using intraluminal impedance. Journal of Pediatrics, 2003;111(4):355-359
- 20.Laranjeira M et al. Adequado crescimento, ausência de ganho de peso excessivo e de alteração no padrão evacuatório em lactentes com refluxo gastroesofágico recebendo fórmula infantil espessada com goma jataí. Pediatr. Mod. 2014; 50(8): 339-342.
- 21.Simakachorn N et al. Randomized, double-blind clinical trial of a lactose-free and a lactose containing formula in dietary management of acute childhood diarrhea. J Med Assoc Thai. 2004 Jun;87(6):641- 9.
- 22.Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. Arq Asma Alerg Imunol. 2018;2(1):7-82.
- 23.Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.
- 24.Niggemann B, Binder C, Dupont C, Hadji S, Arvola T, Isolauri E. Prospective, controlled, multi-center study on the effect of an amino-acid-based formula in infants

with cow's milk allergy/intolerance and atopic dermatitis. *Pediatr Allergy Immunol.* 2001;12(2):78-82.

25.Sampson et al (1992). Safety of an amino acid derived infant formula in children allergic to cow milk. *Pediatrics*, 90(3), 463 465.

26.Verwimp JJ, et al. Symptomatology and growth in infants with cow's milk protein intolerance using two different whey protein hydrolysate based formulas in a Primary Health Care setting. *Eur J Clin Nutr.* 1995;49 Suppl 1:S39 S48.

27.Giampietro PG, et al. Hypoallergenicity of an extensively hydrolyzed whey formula. *Pediatr Allergy Immunol.* 2001;12(2):83 86.

28.Isolauri, E., et al (1995). Efficacy and safety of hydrolyzed cow milk and amino acid derived formulas in infants with cow milk allergy. *The Journal of pediatrics*, 127(4), 550 557.0.

29.Koletzko, Berthold, et al. "Scientific basis and practical application of nutritional care for preterm infants." *Nutritional Care of Preterm Infants*. Vol. 122. Karger Publishers, 2021.

30.Agostoni C et al. Enteral Nutrient Supply for Preterm Infants: Commentary from the European Society for Paediatric Gastroenterology, Hepatology, and Nutrition Committee on Nutrition. *Journal of Pediatric Gastroenterology and Nutrition.* 2010; 50 (1): 85-91