

REFERÊNCIAS APTAMIL E APTANUTRI

1. Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4^a ed. Rio de Janeiro, 2018.
2. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am*. 2013;60(1):49-74.
3. Koletzko B et al. Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *Am J Clin Nutr*. 2009; 89: 1836–45.
4. Dallas DC, Murray NM, Gan J. Proteolytic Systems in Milk: Perspectives on the Evolutionary Function within the Mammary Gland and the Infant. *J Mammary Gland Biol Neoplasia*. 2015 Dec;20(3-4):133-47 .
5. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
6. Liu L et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. *J Lipid Res*. 2014;55(3):531-9
7. Bar-Yoseph F et al. Review of sn-2 palmitate oil implications for infant health. *Prostaglandins Leukot Essent Fatty Acids*. 2013;89(4):139-43
8. Salminen S, Szajewska H, Knol J. The Biotics Family in Early Life. Edited Wiley: vol. 4, 2019
9. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr*. 2002;34(3):291-5
10. Bruzzese E et al. A formula containing galacto- and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study. *Clin Nutr*. 2009;28(2):156-61
11. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr*. 2013;98(2):561S-71S.
12. Scholtens PA et al. Stool characteristics of infants receiving short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides: a review. *World J Gastroenterol*. 2014;20(37):13446-52.
13. Vandenplas Y et al. A partly fermented infant formula with postbiotics including 3'-GL, specific oligosaccharides, 2'-FL, and milk fat supports adequate growth, is safe and well-tolerated in healthy term infants: a double-blind, randomised, controlled, multi-country trial. *Nutrients*. 2020; 12(11):3960
14. Overbeek S et al. Human milk oligosaccharide 2'-fucosyllactose more efficiently modulates immunogenicity during maturation of human dendritic cells in the presence of scGOS/lcFOS prebiotics. *JPGN*. 2019;58:995
15. Comparativo realizado entre produtos da mesma categoria, em 02/2025, utilizando informações extraídas da tabela nutricional
16. Dinleyici M, Barbieur J, Dinleyici EC, Vandenplas Y. Functional effects of human milk oligosaccharides (HMOs). *Gut Microbes*. 2023 Jan-Dec;15(1):2186115.

17. Koletzko, Berthold, et al. "Scientific basis and practical application of nutritional care for preterm infants." Nutritional Care of Preterm Infants. Vol. 122. Karger Publishers, 2021.
18. Embleton, N. D., Moltu, S. J., Lapillonne et al. Enteral nutrition in preterm infants (2022): a position paper from the ESPGHAN committee on nutrition and invited experts. Journal of pediatric gastroenterology and nutrition. 2023; 76(2), 248-268.
19. Aggett, P. J. Agostoni, C. Axelsson I. et al. 2006 Feeding preterm infants after hospital discharge: a commentary by the ESPGHAN Committee on Nutrition. Journal of Pediatric Gastroenterology and Nutrition, 42, 596-603.
20. Lapillonne A, Bronsky J, Campoy C, et al. Feeding the Late and Moderately Preterm Infant: A Position Paper of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. J Pediatr Gastroenterol Nutr. 2019;69(2):259-270.
21. Roy CC et al. Correction of the malabsorption of the preterm infant with a medium-chain triglyceride formula. J Pediatr. 1975;86(3):446-50.
22. Knol J et al. Increase of faecal bifidobacteria due to dietary oligosaccharides induces a reduction of clinically relevant pathogen germs in the faeces of formula-fed preterm infants. Acta Pædiatrica, 2005; 94(Suppl 449): 31–33.
23. Bruzzese E et al. Early administration of GOS/FOS prevents intestinal and respiratory infections in infants. J Pediatr Gastroenterol Nutr. 2006; 42 (5): E95.

REFERÊNCIAS MILNUTRI

1. Nogueira-de-Almeida, C. A., Falcão, M. C., Ribas Filho, D., Zorzo, R. A., Konstantyner, T., Ricci, R., Gioia, N., & Fisberg, M. (2020). Consensus of the Brazilian Association of Nutrology on Milky Feeding of Children Aged 1–5 Years Old. *International Journal of Nutrology*, 13(1), 2–16.
2. Luque et al. Early Programming by Protein Intake: The Effect of Protein on Adiposity Development and the Growth and Functionality of Vital Organs. *Nutrition and Metabolic Insights* 2015;8(S1) 49–56
3. Institute of Medicine (IOM). *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*, 2006.
4. Sociedade Brasileira de Pediatria. *Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola*. Departamento de Nutrologia, 4ª ed. Rio de Janeiro, 2018.
5. Nogueira-de-Almeida, C. A., Vilanova, K. C. M., Perini, T. M., Filho, D. R. (2024). *Alimentação da Criança de Zero a Cinco Anos*. Manual do Departamento de Nutrologia, ABRAN.
6. FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS. *Fats and fatty acids in human nutrition: Report of an expert consultation*. Rome: FAO, 2010.
7. Nogueira-de-Almeida CA, Ribas Filho D, Mello ED, Bertolucci PHF, Falcão MC. I Consenso da Associação Brasileira de Nutrologia sobre recomendações de DHA durante gestação, lactação e infância. In: *Anais do XVIII Congresso Brasileiro de Nutrologia*; 2014; São Paulo, BR. São Paulo;; 2014. 1-13." ; 33.
8. Koletzko, B. (Berthold), editor. *Pediatric Nutrition in Practice*. 2nd, revised edition. (Il. Series: World review of nutrition and dietetics ; vol. 113); 2015
9. Cummings JH, Macfarlane GT. Gastrointestinal effects of prebiotics. *Br J Nutr*. 2002; 87 (Suppl 2):S145-S151.;
10. Veereman G. Pediatric applications of inulin and oligofructose. *J Nutr*. 2007; 137:2585S-9S.
11. Mello, Carolina Santos, Karina Vieira Barros, and Mauro Batista de Moraes, "Brazilian infant and preschool children feeding; literature review." *Jornal de pediatria* 92 (2016): 451-463.
12. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr*. 2002;34(3):291-5.
13. Bruzzese, Eugenia, et al. "A formula containing galacto-and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study." *Clinical Nutrition* 28.2 (2009): 156-161.
14. Shahremian, iraj, et al. "The effects of prebiotic supplementation on weight gain, diarrhoea, constipation, fever and respiratory tract infections in the first year of life" *Journal of paediatrics and child health* 54.8 (2018): 875-880.

15. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. BRASPEN J 2020; 35 (3): 294-306.
16. Liu L et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. J Lipid Res. 2014;55(3):531-9
17. Graf et al. Age dependent incorporation of 14C-DHA into rat brain and body tissues after dosing various 14C-DHA-esters. Prostaglandins, Leukotrienes and Essential Fatty Acids (PLEFA). 2010; 83 (2):89-96.
18. Wijendran V et al. Efficacy of dietary arachidonic acid provided as triglyceride or phospholipid as substrates for brain arachidonic acid accretion in baboon neonates. Pediatr Res. 2002;51(3):265-72
19. Salminen S, Szajewska H, Knol J. The Biotics Family in Early Life. Edited Wiley: vol. 4, 2019
20. Knol J et al. Colon microflora in infants fed formula with galacto- and fructo-oligosaccharides: more like breast-fed infants. J Pediatr Gastroenterol Nutr. 2005.40():36-42.

REFERÊNCIAS FORTINI COMPLETE, FORTINI PLUS, FORTINI PLUS MF, INFATRINI

1. Kerzner B, Milano K, MacLEAn WC, et al. A Practical approach to classifying and managing feeding difficulties. 2015; 135(2):344-35
2. Nogueira CA, Mello E, Filho DR, Maximino P, Fisberg M. Consenso da Associação Brasileira de Nutrologia sobre o uso de suplementos alimentares para crianças com dificuldades alimentares. Int J Nutrol 2018;11(supl S1):S4–S15
3. Sociedade Brasileira de Pediatria. Guia de orientações - Dificuldades alimentares. Departamento Científico de Nutrologia São Paulo: SBP, 2022.
4. Manual de Suporte Nutricional da Sociedade Brasileira de Pediatria - 2ed. - Rio de Janeiro: Departamento Científico de Suporte Nutricional da Sociedade Brasileira de Pediatria. – 2020.
5. Braegger C, Decsi T, Dias JA, Hartman C, Kolaček S, Koletzko B et al. Practical Approach to Paediatric Enteral Nutrition: A Comment by the ESPGHAN Committee on Nutrition. JPGN; 2010;51(1):110-122
6. NICE Guideline. Faltering growth: recognition and management of faltering growth in children. 2017.
7. Mehta NM et al. Defining pediatric malnutrition: A paradigm shift toward etiology-related definitions. J Parenter Enter Nutr. 2013;37(4):460–81.
8. Nogueira-de-Almeida, C.A.; Del Ciampo, L.A.; Martinez, E.Z.; Contini, A.A.; Nogueira-de-Almeida, M.E.; Ferraz, I.S.; Epifanio, M.; da Veiga Ued, F. Clinical Evolution of Preschool Picky Eater Children Receiving Oral Nutritional Supplementation during Six Months: A Prospective Controlled Clinical Trial. Children 2023, 10, 495.
9. Devaera, Y., Syaharutsa, D. M., Jatmiko, H. K., & Sjarif, D. R. Comparing Compliance and Efficacy of Isocaloric Oral Nutritional Supplementation Using 1.5 kcal/mL or 1 kcal/mL Sip Feeds in Mildly to Moderately Malnourished Indonesian Children: A Randomized Controlled Trial. Pediatric gastroenterology, hepatology & nutrition, 21(4), 315–320.
10. Codex Alimentarius FAO/OMS. Standard for infant formula and formulas for special medical purposes intended for infants. Codex Stan 72 – 1981. Revised 2007/2011.
11. Gahagan S. Failure to thrive: a consequence of undernutrition. Pediatr Rev. 2006 Jan;27(1):e1-11.
12. Pencharz PB. Protein and energy requirements for “optimal” catch-up growth. EJCN.2010;64: S5-7
13. Clarke SE, Evans S, MacDonald A, Davies P, Booth IW. Randomized comparison of a nutrient-dense formula with an energy-supplemented formula for infants with faltering growth. J Hum Nutr Diet. 2007;20(4):329-339. 4
14. Comparado a fórmula infantil para lactentes e seu seguimento para lactentes e crianças de primeira infância. Agosto/2024
15. FAO/WHO/UNU. Protein and amino acid requirements in human nutrition: report of a joint FAO/WHO/UNU expert consultation. WHO technical report series no. 935, 2007.p.187

16. Falcão MC, Udsen N, Zamberlan P, Ceccon MER. Rev Bras Nutr Clin 2009; 24 (2): 125-30
17. Uauy R, Hoffman DR, Mena P, Llanos A, Birch EE. Term infant studies of DHA and ARA supplementation on neurodevelopment: results of randomized controlled trials. J Pediatr. 2003;143:S17-25.
18. SanGiovanni JP, Berkey CS, Dwyer JT, Colditz GA. Dietary essential fatty acids, long-chain polyunsaturated fatty acids, and visual resolution acuity in healthy full term infants: a systematic review. Early Hum Dev. 2000;57:165-88.
19. Knol J et al. Colon microflora in infants fed formula with galacto- and fructo-oligosaccharides: more like breast-fed infants. J Pediatr Gastroenterol Nutr. 2005;40(1):36-42.
20. Scholtens PA, Alles MS, Bindels JG, van der Linde EG, Tolboom JJ, Knol J. Bifidogenic effects of solid weaning foods with added prebiotic oligosaccharides: a randomised controlled clinical trial. J Pediatr Gastroenterol Nutr. 2006 May;42(5):553-9.
21. Food and Agriculture Organization of the United Nations. (2010). Fats and fatty acids in human nutrition. Report of an expert consultation. Food and Nutrition Paper 91.
22. Guideline: Sugars intake for adults and children. Geneva: World Health Organization; 2015.
23. Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4ª ed. Rio de Janeiro, 2018.
24. Nogueira-de-Almeida CA, Ribas Filho D, Mello ED, Bertolucci PHF, Falcão MC. I Consenso da Associação Brasileira de Nutrologia sobre recomendações de DHA durante gestação, lactação e infância. In: Anais do XVIII Congresso Brasileiro de Nutrologia; 2014; São Paulo, BR. São Paulo; 2014. 1-13." ; 33.
25. Comparativo realizado entre produtos da mesma categoria, em 02/2025, utilizando informações extraídas da tabela nutricional
26. Scholtens PA et al. Stool characteristics of infants receiving short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides: a review. World J Gastroenterol. 2014;20(37):13446-52.
27. Knol J et al. Colon microflora in infants fed formula with galacto- and fructo-oligosaccharides: more like breast-fed infants. J Pediatr Gastroenterol Nutr. 2005.40():36-42.
28. ASPEN. Nutrition Management of Term Infants with Growth Failure. 2022

REFERÊNCIAS DIAGNÓSTICO E TRATAMENTO DE LACTENTES COM APLV

1. Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. *Arq Asma Alerg Imunol.* 2018;2(1):7-82.
2. Koletzko S et al. Diagnostic approach and management of cow's-milk protein allergy in infants and children: ESPGHAN GI Committee practical guidelines. *J Pediatr Gastroenterol Nutr.* 2012;55 (2):221-9.
3. Muraro A et al. EAACI food allergy and anaphylaxis guidelines: diagnosis and management of food allergy. *Allergy.* 2014;69(8):1008-25.
4. Fiocchi A et al. World Allergy Organization (WAO) Diagnosis and Rationale for Action against Cow's Milk Allergy (DRACMA) Guidelines. *Pediatr Allergy Immunol.* 2010;21 Suppl 21:1-125.
5. Venter C et al. Diagnosis and management of non-IgE-mediated cow's milk allergy in infancy: a UK primary care practical guide. *Clin Transl Allergy.* 2013;3(1):23.
6. Venter C et al. Better recognition, diagnosis and management of non-IgE-mediated cow's milk allergy in infancy: iMAP-an international interpretation of the MAP (Milk Allergy in Primary Care) guideline. *Clin Transl Allergy.* 2017;7:26.
7. Orsi M et al. Alergia a la proteína de la leche de vaca: Propuesta de Guía para el manejo de los niños con alergia a la proteína de la leche de vaca. *Arch. argent. pediatr.* 2009; 107(5): 459-467.
8. Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.
9. Toca MC et al. Consenso sobre el diagnóstico y el tratamiento de la alergia a las proteínas de la leche de vaca de la Sociedad Latinoamericana de Gastroenterología, Hepatología y Nutrición. *Rev Gastroenterol Méx.* 2022;87:235---250.
10. Fox A, Brown T, Walsh J, et al. An update to the Milk Allergy in Primary Care guideline. *Clin Transl Allergy.* 2019;9:40. Published 2019 Aug 12.
11. Vandenplas Y, et al. An ESPGHAN position paper on the diagnosis, management and prevention of cow's milk allergy. *J Pediatr Gastroenterol Nutr.* 2023 Jul 26.
12. Morais MB, Spolidoro JV, Vieira MC, Cardoso AL, Clark O, Nishikawa A, Castro AP. Amino acid formula as a new strategy for diagnosing cow's milk allergy in infants: is it cost-effective? *J Med Econ.* 2016;19:1207-14.
13. Birch E. et al. The DIAMOND (DHA intake and measurement of neural development) study: a double-masked, randomized controlled clinical trial of the maturation of infant visual acuity as a function of the dietary level of docosahexaenoic acid' *Am J Clin Nutr.*, vol. 91, no.4, pg. 848–859, 2010.
14. Qawasmi A et al. Meta-analysis of LCPUFA supplementation of infant formula and visual acuity. *Pediatrics.* 2013;131(1):262-72.

15. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
16. Miles EA, Childs CE, Calder PC. Long-Chain Polyunsaturated Fatty Acids (LCPUFAs) and the Developing Immune System: A Narrative Review. *Nutrients*. 2021; 13: 247.
17. Marten B et al. Medium-chain triglycerides. *International Dairy Journal*. 2006;16: 1374-1382
18. Estudo observacional, prospectivo aberto e multicêntrico de vigilância pós-comercialização sobre aceitabilidade e tolerabilidade de uma fórmula infantil extensamente hidrolisada. Brasil, 2016: dados preliminares.
19. Epifanio M et al. Palatability and Sensory Perception of Infant Formulas for the Treatment of Cow's Milk Allergy According to Brazilian Mothers. *Biomed J Sci & Tech Res* 25(1)-2020. BJSTR. MS.ID.00416.
20. *Nutrologia Pediátrica: Temas da Atualidade em Nutrologia Pediátrica - 2021.* / Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia. São Paulo: SBP, 2021.
21. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am*.2013;60(1):49-74.
22. FERREIRA, CHT; BARRETO, BAP; TOFOLI, MHC; et al. *Microbioma e pediatria na saúde e na doença*. Barueri [SP]: Manole, 2024. 1.ed.
23. *Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar/ Sociedade Brasileira de Pediatria.* – 5.ed. rev. ampl.– São Paulo: SBP, 2024. 208 p.
24. Arslanoglu S, Moro GE, Boehm G, Wienz F, Stahl B, Bertino E. Early neutral prebiotic oligosaccharide supplementation reduces the incidence of some allergic manifestations in the first 5 years of life. *J Biol Regul Homeost Agents*. 2012;26(3 Suppl):49-59.
25. Arslanoglu S, Moro GE, Schmitt J, Tandoi L, Rizzardi S, Boehm G. Early Dietary intervention with a mixture of prebiotic oligosaccharides reduces the incidence of allergic manifestations and infections during the first two years of life. *J Nutr*. 2008;138(6):1091-5.
26. Boehm G. A mixture of prebiotics oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Arch Dis Child*. 2006;91(10):814-
27. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr*. 2002;34(3):291-5
28. Koletzko B, Baker S, Cleghorn G, et al. Global standard for the composition of infant formula: recommendations of an ESPGHAN coordinated international expert group. *J Pediatr Gastroenterol Nutr*. 2005;41(5):584-99.
29. SALMINEN, Seppo; SZAJEWSKA, Hania; KNOL, Jan. *The Biotics Family in Early Life*. 2. ed. Wiley, 2023.

30. Goh CY, et al. Combination of short-chain GOS and long-chain FOS 9:1 with 2FL positively impact the infant gut microbiota composition and metabolic activity in a stimulator of the human intestinal microbial ecosystem (SHIME). *J Pediatr Gastroenterol Nutr* 2019; 68 (S1): NP 114; 1159.
31. Eveleens, R D et al. Definitions, predictors, and outcomes of feeding intolerance in critically ill children: A systematic review. *Clin Nutr*, 2020, 39.3: 685-693.
32. Marino, LV et al. Feeding intolerance in children with critical illness. *Clin. Nutr.*, 2020, 39.3: 609-611.
33. Lucia Diaferio L et al. May Failure to Thrive in Infants Be a Clinical Marker for the Early Diagnosis of Cow's Milk Allergy? *Nutrients* . 2020 Feb 13;12(2):466. doi: 10.3390/nu12020466.
34. Boaventura RM, Mendonça RB, Fonseca FA, Mallozi M, Souza FS, Sarni ROS. Nutritional status and food intake of children with cow's milk allergy. *Allergol Immunopathol (Madr)*. 2019;47(6):544-550. doi:10.1016/j.aller.2019.03.003
35. Meyer R, Venter C, Fox AT, Shah N. Practical dietary management of protein energy malnutrition in young children with cow's milk protein allergy. *Pediatr Allergy Immunol*. 2012;23(4):307-314.

REFERÊNCIAS TRATAMENTO DE CRIANÇAS COM APLV

1. SBP. Guia de orientações - Dificuldades alimentares/ Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia . São Paulo: SBP, 2022.
2. Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. Arq Asma Alerg Imunol. 2018;2(1):7-82.
3. Medeiros LCS et al. Ingestão de nutrientes e estado nutricional de crianças em dieta isenta de leite de vaca e derivados. J. Pediatr. (Rio J.) vol.80 no.5 Porto Alegre, 2004
4. Meyer R, Venter C, Fox AT, Shah N. Practical dietary management of protein energy malnutrition in young children with cow's milk protein allergy. *Pediatr Allergy Immunol* 2012; 23: 307–314.
5. Meyer R et al. A practical approach to vitamin and mineral supplementation in food allergic children. *Clinical and Translational Allergy* (2015) 5:11
6. Tali Sinai et al. Reduced Final Height and Inadequate Nutritional Intake in Cow's Milk-Allergic Young Adults. *J Allergy Clin Immunol Pract* 2019;7:509-15.
7. Maslin K, et al. Comparison of nutrient intake in adolescents and adults with and without food allergies. *J Hum Nutr Diet.* 2018;31(2):209-217.
8. Saad, SMI. Probióticos e prebióticos: o estado de arte. *Rev. Bras. Cienc. Farm.* vol.42 no.1 São Paulo Jan./Mar. 2006.
9. Cummings JH, Macfarlane GT. Gastrointestinal effects of prebiotics. *Br J Nutr.* 2002; 87 (Suppl 2):S145-S151.
10. Veereman G. Pediatric applications of inulin and oligofructose. *J Nutr.* 2007; 137:2585S-9S.
11. Comparativo de tabela nutricional de produtos da mesma categoria no mercado realizada em Agosto 2024.
12. Meyer R, Venter C, Fox AT, Shah N. Practical dietary management of protein energy malnutrition in young children with cow's milk protein allergy. *Pediatr Allergy Immunol.* 2012;23(4):307–14.
13. Joosten K, Meyer R. Nutritional screening and guidelines for managing the child with faltering growth. *Eur J Clin Nutr.* 2010;64:S22–4.
14. WHO/FAO/UNU Expert Consultation. Protein and amino acid requirements in human nutrition: report of a joint. *WHO Tech Rep Ser* [Internet]. 2007;1–265.
15. Lucia Diaferio L et al. May Failure to Thrive in Infants Be a Clinical Marker for the Early Diagnosis of Cow's Milk Allergy?. *Nutrients.* 2020 Feb 13;12(2):466. doi: 10.3390/nu12020466.
16. Boaventura RM, Mendonça RB, Fonseca FA, Mallozi M, Souza FS, Sarni ROS. Nutritional status and food intake of children with cow's milk allergy. *Allergol Immunopathol (Madr).* 2019;47(6):544-550. doi:10.1016/j.aller.2019.03.003
17. Eveleens, R D et al. Definitions, predictors, and outcomes of feeding intolerance in critically ill children: A systematic review. *Clin Nutr,* 2020, 39.3: 685-693.

18. Marino, LV et al. Feeding intolerance in children with critical illness. Clin. Nutr., 2020, 39.3: 609-611.
19. Birch E. et al. The DIAMOND (DHA intake and measurement of neural development) study: a double-masked, randomized controlled clinical trial of the maturation of infant visual acuity as a function of the dietary level of docosahexaenoic acid' Am J Clin Nutr., vol. 91, no.4, pg. 848–859, 2010.
20. Qawasmi A et al. Meta-analysis of LCPUFA supplementation of infant formula and visual acuity. Pediatrics. 2013;131(1):262-72.
21. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. BRASPEN J 2020; 35 (3): 294-306
22. Miles EA, Childs CE, Calder PC. Long-Chain Polyunsaturated Fatty Acids (LCPUFAs) and the Developing Immune System: A Narrative Review. Nutrients. 2021; 13: 247.
23. Mabin DC et al. Nutritional content of few foods diet in atopic dermatitis. Arch Dis Child. 1995;73(3):208-10.
24. Marten B et al. Medium-chain triglycerides. International Dairy Journal. 2006;16: 1374-1382

REFERÊNCIAS LINHA SOJA

1. Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. Arq Asma Alerg Imunol. 2018;2(1):7-82.
2. Bhatia J, Greer F. American Academy of Pediatrics Committee on Nutrition: Use of soy protein-based formulas in infant feeding. Pediatrics. 2008;121(5):1062-8
3. Agostoni C et al. Soy protein infant formulae and follow-on formulae: A commentary by the ESPGHAN Committee on Nutrition. J Ped Gast Nutr 2006: 42:352-61.
4. Vandenplas Y, Hegar B, Munasir Z, et al. The role of soy plant-based formula supplemented with dietary fiber to support children's growth and development: An expert opinion. Nutrition. 2021;90:111278.
5. Merritt RJ, Fleet SE, Fifi A, et al. North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition Position Paper: Plant-based Milks. J Pediatr Gastroenterol Nutr. 2020;71(2):276-281.
6. ILSI (International Life Sciences Institute) Brasil. Série: Funções plenamente reconhecidas de Nutrientes. 1 Ed. 2009-2014.

REFERÊNCIAS LINHA GASTRO

1. Rosen R, Vandenplas Y, Singendonk M, et al. Pediatric Gastroesophageal Reflux Clinical Practice Guidelines: Joint Recommendations of the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition and the European Society for Pediatric Gastroenterology, Hepatology, and Nutrition. *J Pediatr Gastroenterol Nutr.* 2018;66(3):516-554.
2. Salvatore, S. et al. Thickened infant formula: what to know. *Nutrition*, 2018.
3. Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation, and cow's milk allergy in formula-fed infants. *Acta Pædiatrica* 2015,
4. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5.
5. Braegger C et al. Supplementation of infant formula with probiotics and/or prebiotics: a systematic review and comment by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr.* 2011;52:238–225.
6. Boehm G et al. Prebiotics in infant formulas. *J Clin Gastroenterol.* 2004; 38: S76-9.
7. Vandenplas Y et al. Oligosaccharides in infant formula: more evidence to validate the role of prebiotics. *Br J Nutr.* 2015;113(9):1339-44.
8. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr.* 2013;98(2):561S-71S.
9. Infante Pina D et al. Thickened infant formula, rheological study of the “in vitro” properties. *An Pediatr (Barc).* 2010; 72(5):302-8.
10. Horvath A, Dziechciarz P, Szajewska H. The Effect of Thickened-Feed Interventions on Gastroesophageal Reflux in Infants: Systematic Review and Meta-analysis of Randomized, Controlled Trials. *Pediatrics* 2008;122:e1268–e1277.
11. Koletzko B, Karin B, Thomas BJ, Phipip C Should formula for infants provide arachidonic acid along with DHA? A position paper of the European Academy of Paediatrics and the Child Health Foundation. *Am J Clin Nutri.* 2020. Vol. 111, nº 1, p. 10-16.
12. Birch E. et al. The DIAMOND (DHA intake and measurement of neural development) study: a double-masked, randomized controlled clinical trial of the maturation of infant visual acuity as a function of the dietary level of docosahexaenoic acid' *Am J Clin Nutr.*, vol. 91, no.4, pg. 848–859, 2010.
13. Qawasmi A et al. Meta-analysis of LCPUFA supplementation of infant formula and visual acuity. *Pediatrics.* 2013;131(1):262-72.
14. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
15. Miles EA, Childs CE, Calder PC. Long-Chain Polyunsaturated Fatty Acids (LCPUFAs) and the Developing Immune System: A Narrative Review. *Nutrients.* 2021; 13: 247.
16. Lerner A, Shamir R. Nucleotides in infant nutrition: a must or an option. *Isr Med Assoc J.* 2000; 2(10):772-4.

17. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am.* 2013;60(1):49-74.
18. Dupont C. Protein requirements during the first year of life. *Am J Clin Nutr.* 2003;77(6):1544S-1549S.
19. EFSA. Scientific Opinion on the essential composition of infant and follow-on formulae. *EFSA Journal.* 2014;12(7).
20. Sociedade Brasileira de Pediatria. Departamento de Nutrologia. Manual de Alimentação: Da Infância à adolescência, 2018
21. Borrelli O et al. Use of a new thickened formula for treatment of symptomatic gastroesophageal reflux in infants. *Ital J Gastroenterol Hepatol.* 1997; 29:237-42.
22. Wenzl TG et al. Effects of thickened feeding on gastroesophageal reflux in infants: a placebo-controlled crossover study using intraluminal impedance. *Journal of Pediatrics.* 2003;111(4):355-359.
23. Danone Nutricia Research: <https://www.nutriciaaresearch.com>.
24. Meunier L et al. Locust bean gum safety in neonates and young infants: an integrated review of the toxicological database and clinical evidence. *Regul Toxicol Pharmacol.* 2014;70(1):155-69.
25. Laranjeira M et al. Adequado crescimento, ausência de ganho de peso excessivo e de alteração no padrão evacuatório em lactentes com refluxo gastroesofágico recebendo fórmula infantil espessada com goma jataí. *Pediatr. Mod.* 2014; 50(8): 339-342.
26. Savino F et al. Reduction of crying episodes owing to infantile colic: A randomized controlled study on the efficacy of a new infant formula. *Eur J Clin Nutr* 2006, 60:1304-10.
27. Savino F et al. "Minor" feeding problems during the first months of life: effect of a partially hydrolyzed milk formula containing fructo- and galacto-oligosaccharides. *Acta Paediatr Suppl* 2003, 91:86-90.
28. Schmelzle H et al. Randomized double-blind study of the nutritional efficacy and bifidogenicity of a new infant formula containing partially hydrolyzed protein, a high beta-palmitic acid level, and nondigestible oligosaccharides. *J Pediatr Gastroenterol Nutr* 2003, 36:343-51.
29. Bongers ME et al. The clinical effect of a new infant formula in term infants with constipation: a double-blind, randomized cross-over trial. *Nutr J* 2007;6:8.
30. Savino F et al. Advances in the management of digestive problems during the first months of life. *Acta Paediatr Suppl* 2005, 94:120.
31. Havlicekova Z et al. Beta-palmitate - a natural component of human milk in supplemental milk formulas. *Nutr J.* 2016;15(1):28.
32. Litmanovitz I et al. Reduced crying in term infants fed high beta-palmitate formula: a double-blind randomized clinical trial. *BMC Pediatr.* 2014;14:152.
33. Yao M et al. Effects of term infant formulas containing high sn-2 palmitate with and without oligofructose on stool composition, stool characteristics, and bifidogenicity. *J Pediatr Gastroenterol Nutr.* 2014 ;59(4):440-8.
34. Nowacki J et al. Stool fatty acid soaps, stool consistency and gastrointestinal tolerance in term infants fed infant formulas containing

- high sn-2 palmitate with or without oligofructose: a double-blind, randomized clinical trial. *Nutr J.* 2014;13:105.
35. Simakachorn N et al. Randomized, double-blind clinical trial of a lactose-free and a lactose containing formula in dietary management of acute childhood diarrhea. *J Med Assoc Thai.* 2004 Jun;87(6):641- 9.
 36. Comparativo de tabela nutricional de produtos da mesma categoria no mercado realizada em Agosto 2024.