

REFERÊNCIAS

ONE PED DANONE

1. Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4^a ed. Rio de Janeiro, 2018.
2. Zubler et al. Evidence-informed milestones for developmental surveillance tools Pediatrics. 2022;149 (3)
3. Nogueira-de-Almeida, C. A., Falcão, M. C., Ribas Filho, D., Zorzo, R. A., Konstantyner, T., Ricci, R., Gioia, N., & Fisberg, M. (2022). Consensus of the Brazilian Association of Nutrology on Milky Feeding of Children Aged 1–5 Years Old. International Journal of Nutrology, 13(1), 2–16.
4. Manual de Suporte Nutricional da Sociedade Brasileira de Pediatria - 2ed. - Rio de Janeiro: Departamento Científico de Suporte Nutricional da Sociedade Brasileira de Pediatria. – 2020.
5. SBV. Pesquisa do IBOPE aponta crescimento histórico no número de vegetarianos no Brasil, Disponível em: Pesquisa do IBOPE aponta crescimento histórico no número de vegetarianos no Brasil - SVB. Acesso em 01.03.24.
6. Savilahti et al. 2012. 29.
7. Merritt, R. et al. North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition Position Paper: Plantbased Milks. JPGN Volume 71, Number 2, August 2020.
8. Sociedade Brasileira de Pediatria. Guia de orientações - Dificuldades alimentares. Departamento Científico de Nutrologia São Paulo: SBP, 2022.
9. Codex Alimentarius FAO/OMS. Standard for infant formula and formulas for special medical purposes intended for infants. Codex Stan 72 – 1981. Revised 2007/2011.
10. Gahagan S. Failure to thrive: a consequence of undernutrition. Pediatr Rev. 2006 Jan;27(1):e1-11.
11. Benninga MA, Faure C, Hyman PE, et al. Childhood functional gastrointestinal disorders:neonate/toddler. Gastroenterology 2016;150:1443-55.
12. Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation and cow's milk allergy in formula-fed infants. Acta Paediatrica 2015, ISSN 0803-5253
13. Savino F et al. Reduction of crying episodes owing to infantile colic: A randomized controlled study on the efficacy of a new infant formula. Eur J Clin Nutr 2006, 60:1304-10.
14. Savino F et al. "Minor" feeding problems during the first months of life: effect of a partially hydrolyzed milk formula containing fructo- and galacto-oligosaccharides. Acta Paediatr Suppl 2003, 91:86-90.
15. Schmelzle H et al. Randomized double-blind study of the nutritional efficacy and bifidogenicity of a new infant formula containing partially hydrolyzed protein, a high beta-palmitic acid level, and nondigestible oligosaccharides. J Pediatr Gastroenterol Nutr 2003, 36:343-51.
16. Bongers ME et al. The clinical effect of a new infant formula in term infants with constipation: a double-blind, randomized cross-over trial. Nutr J 2007;6:8.
17. Savino F et al. Advances in the management of digestive problems during the first months of life. Acta Paediatr Suppl 2005, 94:120.
18. Borrelli O et al. Use of a new thickened formula for treatment of symptomatic gastroesophageal reflux in infants. Ital J Gastroenterol Hepatol, 1997; 29:237-42.
19. Wenzl TG et al. Effects of thickened feeding on gastroesophageal reflux in infants: a placebo-controlled crossover study using intraluminal impedance. Journal of Pediatrics, 2003;111(4):355-359
20. Laranjeira M et al. Adequado crescimento, ausência de ganho de peso excessivo e de alteração no padrão evacuatório em lactentes com refluxo gastroesofágico recebendo fórmula infantil espessada com goma jataí. Pediatr. Mod. 2014; 50(8): 339-342.

21. Simakachorn N et al. Randomized, double-blind clinical trial of a lactose-free and a lactose containing formula in dietary management of acute childhood diarrhea. *J Med Assoc Thai.* 2004 Jun;87(6):641- 9.
22. Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. *Arq Asma Alerg Imunol.* 2018;2(1):7-82.
23. Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.
24. Niggemann B, Binder C, Dupont C, Hadji S, Arvola T, Isolauri E. Prospective, controlled, multi-center study on the effect of an amino-acid-based formula in infants with cow's milk allergy/intolerance and atopic dermatitis. *Pediatr Allergy Immunol.* 2001;12(2):78-82.
25. Sampson et al (1992). Safety of an amino acid derived infant formula in children allergic to cow milk. *Pediatrics,* 90(3), 463 465.
26. Verwimp JJ, et al. Symptomatology and growth in infants with cow's milk protein intolerance using two different whey protein hydrolysate based formulas in a Primary Health Care setting. *Eur J Clin Nutr.* 1995;49 Suppl 1:S39 S48.
27. Giampietro PG, et al. Hypoallergenicity of an extensively hydrolyzed whey formula. *Pediatr Allergy Immunol.* 2001;12(2):83 86.
28. Isolauri, E., et al (1995). Efficacy and safety of hydrolyzed cow milk and amino acid derived formulas in infants with cow milk allergy. *The Journal of pediatrics,* 127(4), 550 557.0.
29. Koletzko, Berthold, et al. "Scientific basis and practical application of nutritional care for preterm infants." *Nutritional Care of Preterm Infants.* Vol. 122. Karger Publishers, 2021.
30. Agostoni C et al. Enteral Nutrient Supply for Preterm Infants: Commentary from the European Society for Paediatric Gastroenterology, Hepatology, and Nutrition Committee on Nutrition. *Journal of Pediatric Gastroenterology and Nutrition.* 2010; 50 (1): 85-91

REFERÊNCIAS APTAMIL E APTANUTRI

1. Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4^a ed. Rio de Janeiro, 2018.
2. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am.* 2013;60(1):49-74.
3. Koletzko B et al. Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *Am J Clin Nutr.* 2009; 89: 1836–45.
4. Dallas DC, Murray NM, Gan J. Proteolytic Systems in Milk: Perspectives on the Evolutionary Function within the Mammary Gland and the Infant. *J Mammary Gland Biol Neoplasia.* 2015 Dec;20(3-4):133-47 .
5. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
6. Liu L et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. *J Lipid Res.* 2014;55(3):531-9
7. Bar-Yoseph F et al. Review of sn-2 palmitate oil implications for infant health. *Prostaglandins Leukot Essent Fatty Acids.* 2013;89(4):139-43
8. Salminen S, Szajewska H, Knol J. The Biotics Family in Early Life. Edited Wiley: vol. 4, 2019
9. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5
10. Bruzzese E et al. A formula containing galacto- and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study. *Clin Nutr.* 2009;28(2):156-61
11. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr.* 2013;98(2):561S-71S.
12. Scholtens PA et al. Stool characteristics of infants receiving short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides: a review. *World J Gastroenterol.* 2014;20(37):13446-52.

13. Arrieta MC, Stiensma LT, Amenyogbe N, Brown EM, Finlay B. The intestinal microbiome in early life: health and disease. *Front Immunol.* 2014 Sep 5;5:427.
14. Vandenplas Y et al. A partly fermented infant formula with postbiotics including 3'-GL, specific oligosaccharides, 2'-FL, and milk fat supports adequate growth, is safe and well-tolerated in healthy term infants: a double-blind, randomised, controlled, multi-country trial. *Nutrients.* 2020; 12(11):3960
15. Overbeek S et al. Human milk oligosaccharide 2'-fucosyllactose more efficiently modulates immunogenicity during maturation of human dendritic cells in the presence of scGOS/IcFOS prebiotics. *JPGN.* 2019;58:995
16. Comparativo realizado entre produtos da mesma categoria, em 02/2025, utilizando informações extraídas da tabela nutricional
17. Mun JG, Legette LL, Ikonte CJ, Mitmesser SH. Choline and DHA in Maternal and Infant Nutrition: Synergistic Implications in Brain and Eye Health. *Nutrients.* 2019 May 21;11(5):1125.

REFERÊNCIAS FORTINI COMPLETE, FORTINI PLUS, FORTINI PLUS MF, INFATRINI

1. Sociedade Brasileira de Pediatria – Departamento de Nutrologia. Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar - 4ª. ed. - São Paulo: SBP, 2018
2. Nogueira CA, Mello E, Filho DR, Maximino P, Fisberg M. Consenso da Associação Brasileira de Nutrologia sobre o uso de suplementos alimentares para crianças com dificuldades alimentares. *Int J Nutrol* 2018;11(suppl S1):S4–S15
3. Sociedade Brasileira de Pediatria. Guia de orientações - Dificuldades alimentares. Departamento Científico de Nutrologia São Paulo: SBP, 2022.
4. Manual de Suporte Nutricional da Sociedade Brasileira de Pediatria - 2ed. - Rio de Janeiro: Departamento Científico de Suporte Nutricional da Sociedade Brasileira de Pediatria. – 2020.
5. Codex Alimentarius FAO/OMS. Standard for infant formula and formulas for special medical purposes intended for infants. Codex Stan 72 – 1981. Revised 2007/2011.
6. Gahagan S. Failure to thrive: a consequence of undernutrition. *Pediatr Rev.* 2006 Jan;27(1):e1-11.

REFERÊNCIAS LINHA GASTRO

1. Salvatore, S. et al. Thickened infant formula: what to know. *Nutrition*, 2018.
2. Rosen R, Vandenplas Y, Singendonk M, et al. Pediatric Gastroesophageal Reflux Clinical Practice Guidelines: Joint Recommendations of the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition and the European Society for Pediatric Gastroenterology, Hepatology, and Nutrition. *J Pediatr Gastroenterol Nutr.* 2018;66(3):516-554.
3. Infante Pina D et al. Thickened infant formula, rheological study of the “in vitro” properties. *An Pediatr (Barc).* 2010; 72(5):302-8.
4. Horvath A, Dziechciarz P, Szajewska H. The Effect of Thickened-Feed Interventions on Gastroesophageal Reflux in Infants: Systematic Review and Meta-analysis of Randomized, Controlled Trials. *Pediatrics* 2008;122:e1268–e1277.
5. Comparativo de tabela nutricional de produtos da mesma categoria realizado em Abril, 2025.
6. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5
7. Braegger C et al. Supplementation of infant formula with probiotics and/or prebiotics: a systematic review and comment by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr.* 2011;52:238–225
8. Boehm G et al. Prebiotics in infant formulas. *J Clin Gastroenterol.* 2004; 38: S76-9.

- 9.Vandenplas Y et al. Oligosaccharides in infant formula: more evidence to validate the role of prebiotics. *Br J Nutr.* 2015;113(9):1339-44.
- 10.Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr.* 2013;98(2):561S-71S.
- 11.Koletzko B, Karin B, Thomas BJ, Phipip C Should formula for infants provide arachidonic acid along with DHA? A position paper of the European Academy of Paediatrics and the Child Health Foundation. *Am J Clin Nutri.* 2020. Vol. 111, nº 1, p. 10-16.
- 12.Birch E. et al. The DIAMOND (DHA intake and measurement of neural development) study: a double-masked, randomized controlled clinical trial of the maturation of infant visual acuity as a function of the dietary level of docosahexaenoic acid' *Am J Clin Nutr.*, vol. 91, no.4, pg. 848–859, 2010.
- 13.Qawasmi A et al. Meta-analysis of LCPUFA supplementation of infant formula and visual acuity. *Pediatrics.* 2013;131(1):262-72.
- 14.Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
- 15.Miles EA, Childs CE, Calder PC. Long-Chain Polyunsaturated Fatty Acids (LCPUFAs) and the Developing Immune System: A Narrative Review. *Nutrients.* 2021; 13: 247.
16. Lerner A, Shamir R. Nucleotides in infant nutrition: a must or an option. *Isr Med Assoc J.* 2000;2(10):772-4.
- 17.Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am.*2013;60(1):49-74.
18. Danone Nutricia Research: <https://www.nutriciaresearch.com>. Acesso em agosto de 2020.
19. Borrelli O et al. Use of a new thickened formula for treatment of symptomatic gastroesophageal reflux in infants. *Ital J Gastroenterol Hepatol,* 1997; 29:237-42.
20. Wenzl TG et al. Effects of thickened feeding on gastroesophageal reflux in infants: a placebo-controlled crossover study using intraluminal impedance. *Journal of Pediatrics,* 2003;111(4):355-359
21. Rosen R, Vandenplas Y, Singendonk M, et al. Pediatric Gastroesophageal Reflux Clinical Practice Guidelines: Joint Recommendations of the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition and the European Society for Pediatric Gastroenterology, Hepatology, and Nutrition. *J Pediatr Gastroenterol Nutr.* 2018;66(3):516-554.
22. Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation and cow's milk allergy in formula-fed infants. *Acta Pædiatrica* 2015, ISSN 0803-5253