

REFERÊNCIAS BIBLIOGRÁFICAS

One Ped Danone

1. Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4ª ed. Rio de Janeiro, 2018.
2. Zubler et al. Evidence-informed milestones for developmental surveillance tools Pediatrics. 2022;149 (3)
3. Nogueira-de-Almeida, C. A., Falcão, M. C., Ribas Filho, D., Zorzo, R. A., Konstantyner, T., Ricci, R., Gioia, N., & Fisberg, M. (2022). Consensus of the Brazilian Association of Nutrology on Milky Feeding of Children Aged 1–5 Years Old. International Journal of Nutrology, 13(1), 2–16.
4. Manual de Suporte Nutricional da Sociedade Brasileira de Pediatria - 2ed. - Rio de Janeiro: Departamento Científico de Suporte Nutricional da Sociedade Brasileira de Pediatria. – 2020.
5. SBV. Pesquisa do IBOPE aponta crescimento histórico no número de vegetarianos no Brasil, Disponível em: Pesquisa do IBOPE aponta crescimento histórico no número de vegetarianos no Brasil - SVB. Acesso em 01.03.24.
6. Savilahti et al. 2012. 29.
7. Merritt, R. et al. North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition Position Paper: Plantbased Milks. JPGN Volume 71, Number 2, August 2020.
8. Sociedade Brasileira de Pediatria. Guia de orientações - Dificuldades alimentares. Departamento Científico de Nutrologia São Paulo: SBP, 2022.
9. Codex Alimentarius FAO/OMS. Standard for infant formula and formulas for special medical purposes intended for infants. Codex Stan 72 – 1981. Revised 2007/2011.
10. Gahagan S. Failure to thrive: a consequence of undernutrition. Pediatr Rev. 2006 Jan;27(1):e1-11.
11. Benninga MA, Faure C, Hyman PE, et al. Childhood functional gastrointestinal disorders: neonate/toddler. Gastroenterology 2016;150:1443-55.
12. Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation and cow's milk allergy in formula-fed infants. Acta Paediatrica 2015, ISSN 0803-5253
13. Savino F et al. Reduction of crying episodes owing to infantile colic: A randomized controlled study on the efficacy of a new infant formula. Eur J Clin Nutr 2006, 60:1304-10.
14. Savino F et al. "Minor" feeding problems during the first months of life: effect of a partially hydrolyzed milk formula containing fructo- and galacto-oligosaccharides. Acta Paediatr Suppl 2003, 91:86-90.
15. Schmelzle H et al. Randomized double-blind study of the nutritional efficacy and bifidogenicity of a new infant formula containing partially hydrolyzed protein, a high

- beta-palmitic acid level, and nondigestible oligosaccharides. *J Pediatr Gastroenterol Nutr* 2003, 36:343-51.
16. Bongers ME et al. The clinical effect of a new infant formula in term infants with constipation: a double-blind, randomized cross-over trial. *Nutr J* 2007;6:8.
 17. Savino F et al. Advances in the management of digestive problems during the first months of life. *Acta Paediatr Suppl* 2005, 94:120.
 18. Borrelli O et al. Use of a new thickened formula for treatment of symptomatic gastroesophageal reflux in infants. *Ital J Gastroenterol Hepatol*, 1997; 29:237-42.
 19. Wenzl TG et al. Effects of thickened feeding on gastroesophageal reflux in infants: a placebo-controlled crossover study using intraluminal impedance. *Journal of Pediatrics*, 2003;111(4):355-359
 20. Laranjeira M et al. Adequado crescimento, ausência de ganho de peso excessivo e de alteração no padrão evacuatório em lactentes com refluxo gastroesofágico recebendo fórmula infantil espessada com goma jataí. *Pediatr. Mod.* 2014; 50(8): 339-342.
 21. Simakachorn N et al. Randomized, double-blind clinical trial of a lactose-free and a lactose containing formula in dietary management of acute childhood diarrhea. *J Med Assoc Thai.* 2004 Jun;87(6):641- 9.
 22. Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. *Arq Asma Alerg Imunol.* 2018;2(1):7-82.
 23. Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.
 24. Niggemann B, Binder C, Dupont C, Hadji S, Arvola T, Isolauri E. Prospective, controlled, multi-center study on the effect of an amino-acid-based formula in infants with cow's milk allergy/intolerance and atopic dermatitis. *Pediatr Allergy Immunol.* 2001;12(2):78-82.
 25. Sampson et al (1992). Safety of an amino acid derived infant formula in children allergic to cow milk. *Pediatrics*, 90(3), 463 465.
 26. Verwimp JJ, et al. Symptomatology and growth in infants with cow's milk protein intolerance using two different whey protein hydrolysate based formulas in a Primary Health Care setting. *Eur J Clin Nutr.* 1995;49 Suppl 1:S39 S48.
 27. Giampietro PG, et al. Hypoallergenicity of an extensively hydrolyzed whey formula. *Pediatr Allergy Immunol.* 2001;12(2):83 86.
 28. Isolauri, E., et al (1995). Efficacy and safety of hydrolyzed cow milk and amino acid derived formulas in infants with cow milk allergy. *The Journal of pediatrics*, 127(4), 550 557.0.
 29. Koletzko, Berthold, et al. "Scientific basis and practical application of nutritional care for preterm infants." *Nutritional Care of Preterm Infants.* Vol. 122. Karger Publishers, 2021.
 30. Agostoni C et al. Enteral Nutrient Supply for Preterm Infants: Commentary from the European Society for Paediatric Gastroenterology, Hepatology, and Nutrition Committee on Nutrition. *Journal of Pediatric Gastroenterology and Nutrition.* 2010; 50 (1): 85-91

Referências Aptamil

1. Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4ª ed. Rio de Janeiro, 2018.
2. Zubler et al. Evidence-informed milestones for developmental surveillance tools Pediatrics. 2022; 149 (3)
3. Moreno Villares JM, Collado MC, Larqué E, Leis Trabazo R, Saenz De Pipaón M, Moreno Aznar LA. The first 1000 days: an opportunity to reduce the burden of noncommunicable diseases. Nutr Hosp. 2019 Mar 7;36(1):218-232.
4. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. Pediatr Clin North Am. 2013;60(1):49-74.
5. Koletzko B et al. Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. Am J Clin Nutr. 2009; 89: 1836–45.
6. Dallas DC, Murray NM, Gan J. Proteolytic Systems in Milk: Perspectives on the Evolutionary Function within the Mammary Gland and the Infant. J Mammary Gland Biol Neoplasia. 2015 Dec;20(3-4):133-47 .
7. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. BRASPEN J 2020; 35 (3): 294-306
8. Liu L et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. J Lipid Res. 2014;55(3):531-9
9. Bar-Yoseph F et al. Review of sn-2 palmitate oil implications for infant health. Prostaglandins Leukot Essent Fatty Acids. 2013;89(4):139-43
10. Salminen S, Szajewska H, Knol J. The Biotics Family in Early Life. Edited Wiley: vol. 4, 2019
11. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. J Pediatr Gastroenterol Nutr. 2002;34(3):291-5
12. Bruzzese E et al. A formula containing galacto- and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study. Clin Nutr. 2009;28(2):156-61
13. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. Am J Clin Nutr. 2013;98(2):561S-71S.
14. Scholtens PA et al. Stool characteristics of infants receiving short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides: a review. World J Gastroenterol. 2014;20(37):13446-52.

15. Vandenplas Y et al. A partly fermented infant formula with postbiotics including 3'-GL, specific oligosaccharides, 2'-FL, and milk fat supports adequate growth, is safe and well-tolerated in healthy term infants: a double-blind, randomised, controlled, multi-country trial. *Nutrients*. 2020; 12(11):3960
16. Overbeek S et al. Human milk oligosaccharide 2'-fucosyllactose more efficiently modulates immunogenicity during maturation of human dendritic cells in the presence of scGOS/lcFOS prebiotics. *JPGN*. 2019;58:995
17. Comparativo realizado entre produtos da mesma categoria, em 04/2025, utilizando informações extraídas da tabela nutricional
18. Mun JG, Legette LL, Ikente CJ, Mitmesser SH. Choline and DHA in Maternal and Infant Nutrition: Synergistic Implications in Brain and Eye Health. *Nutrients*. 2019 May 21;11(5):1125.

Referências Aptanutri

1. SBP. Sociedade Brasileira de Pediatria – Departamento de Nutrologia Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar / Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia. – 4ª. ed. - São Paulo: SBP, 2018. 172 p
2. Zubler et al. Evidence-informed milestones for developmental surveillance tools *Pediatrics*. 2022; 149 (3).
3. Nogueira-de-Almeida, C. A., Falcão, M. C., Ribas Filho, D., Zorzo, R. A., Konstantyner, T., Ricci, R., Gioia, N., & Fisberg, M. (2022). Consensus of the Brazilian Association of Nutrology on Milky Feeding of Children Aged 1–5 Years Old. *International Journal of Nutrology*, 13(1), 2–16.
4. Nogueira-de-Almeida, C. A., Vilanova, K. C. M., Perini, T. M., Filho, D. R. (2024). Alimentação da Criança de Zero a Cinco Anos. Manual do Departamento de Nutrologia, ABRAN.
5. Hojsak, I. et al. Young Child Formula: A Position Paper by the ESPGHAN Committee on Nutrition *JPGN* 2018;66: 177-185.
6. SBP. Sociedade Brasileira de Pediatria - Departamento Científico de Endocrinologia. Avaliação do crescimento: O quê o pediatra precisa saber, 2023.
7. Dekaban, A; Sadowsky, D. Changes in brain weights during the span of human life: Relation of brain weights to body heights and body weights. *Annals of Neurology* Vol. 4 Issue 4 Oct. 1978
8. Lenroot, R; Giedd, J.N. Brain development in children and adolescents: insights from anatomical magnetic resonance imaging. *Neurosci Biobehav Rev*. 2006;30(6):718-2
9. Bernardi, J.R. et al. Fetal and Neonatal Levels of Omega-3: Effects on Neurodevelopment, Nutrition, and Growth. <https://doi.org/10.1100/2012/202473>
10. Colombo, J. et al. Long Chain Polyunsaturated Fatty Acid Supplementation in Infancy Reduces Heart Rate and Positively Affects Distribution of Attention. *Pediatr Res*. 2011 Oct; 70(4): 406–410.

11. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
12. Eussem SRBM, et al. Theoretical Impact of Replacing Whole Cow's Milks by Young-Child Formula on Nutrient Intakes of UK Young Children: Results of a Simulation Study. *Annals of Nutrition and Metabolism*, 2015; 67(4):247-56)
13. Institute of Medicine. Dietary Reference Intakes; The Essential Guide to Nutrient Requirements (2006) - Comparativo vs as necessidades nutricionais de adultos
14. Mello CS, Barros KV, Moraes MB. Brazilian infant and preschool children feeding: literature review. *J Pediatr(Rio J)*. 2016;92:451---63.
15. Luque et al. Early Programming by Protein Intake: The Effect of Protein on Adiposity Development and the Growth and Functionality of Vital Organs. *Nutrition and Metabolic Insights* 2015;8(S1) 49–56.
16. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am*. 2013;60(1):49-74.
17. Koletzko B et al. Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *Am J Clin Nutr*. 2009; 89: 1836–45.
18. Dallas DC, Murray NM, Gan J. Proteolytic Systems in Milk: Perspectives on the Evolutionary Function within the Mammary Gland and the Infant. *J Mammary Gland Biol Neoplasia*. 2015 Dec;20(3-4):133-47.
19. Arrieta MC, Stiemsma LT, Amenyogbe N, Brown EM, Finlay B. The intestinal microbiome in early life: health and disease. *Front Immunol*. 2014 Sep 5;5:427. doi: 10.3389/fimmu.2014.00427. PMID: 25250028; PMCID: PMC4155789.
20. Salminen S, Szajewska H, Knel J. The Biotics Family in Early Life, Edited Wiley: vol. 4, 2019
21. Dekaban, A; Sadowsky, D. Changes in brain weights during the span of human life: Relation of brain weights to body heights and body weights. *Annals of Neurology* Vol. 4 Issue 4 Oct. 1978.
22. Lenroot, R; Giedd, J.N. Brain development in children and adolescents: insights from anatomical magnetic resonance imaging. *Neurosci Biobehav Rev*. 2006;30(6):718-2.
23. Colombo, J. et al. Long Chain Polyunsaturated Fatty Acid Supplementation in Infancy Reduces Heart Rate and Positively Affects Distribution of Attention. *Pediatr Res*. 2011 Oct; 70(4): 406–410.
24. Costantini L et al: Impact of Omega-3 Fatty Acids on the Gut Microbiota, *Int. J. Mol. Sci*. 2017;18(2), 2645.
25. Gilmore JH, Knickmeyer RC, Gao W. Imaging structural and functional brain development in early childhood. *Nat Rev Neurosci*. 2018 Feb 16;19(3):123-137. doi: 10.1038/nrn.2018.1. PMID: 29449712; PMCID: PMC5987539.
26. Mun JG, Legette LL, Ikente CJ, Mitmesser SH. Choline and DHA in Maternal and Infant Nutrition: Synergistic Implications in Brain and Eye Health. *Nutrients*. 2019 May 21;11(5):1125.
27. Comparativo realizado entre produtos da mesma categoria, em 03/2025, utilizando informações extraídas da tabela nutricional
28. Liu L et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. *J Lipid Res*. 2014;55(3):531-9

29. Graf et al Age dependent incorporation of 14C-DHA into rat brain and body tissues after dosing various 14C-DHA-esters. Prostaglandins, Leukotrienes and Essential Fatty Acids (PLEFA). 2010; 83 (2):89-96.
30. Wijendran V et al. Efficacy of dietary arachidonic acid provided as triglyceride or phospholipid as substrates for brain arachidonic acid accretion in baboon neonates, *Pediatr Res*. 2002;51(43) 265-73
31. Moro G et al. Desago-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr*. 2002;34(3):291-5.
32. Shahremian, iraj, et al. "The effects of prebiotic supplementation on weight gain, diarrhoea, constipation, fever and respiratory tract infections in the first year of life" *Journal of paediatrics and child health* 54.8 (2018): 875-880
33. Kranz, S., Brauchla M., Slavin, J. L., & Miller, K. B. (2012). What do we know about dietary fiber intake in children and health? The effects of fiber intake on constipation, obesity, and diabetes in children. *Advances in Nutrition*, 3(1), 47-53.
34. Bruzzese, Eugenia, et al. "A formula containing galacto-and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study." *Clinical Nutrition* 28.2 (2009): 156-161.
35. Resolução da Diretoria Colegiada - RDC nº 44, de 19 de setembro de 2011
36. UNIVERSIDADE FEDERAL DO RIO DE JANEIRO. Biomarcadores do estado de micronutrientes: prevalências de deficiências e curvas de distribuição de micronutrientes em crianças brasileiras menores de 5 anos 3: ENANI 2019. - Documento eletrônico. - Rio de Janeiro, RJ: UFRJ, 2021. (156 p.). Coordenador geral, Gilberto Kac. Disponível em: <https://enani.nutricao.ufrj.br/index.php/relatorios/>. Acesso em: 12.03.2025
37. Bar-Yoseph F et al, Review of sn-2 palmitate oil implications for infant health *Prostaglandins Leukot Essent Fatty Acids*. 2013;89(4)139-43.
38. Yaron, S. et al. *Nutr*. 2013 Apr 56(43):376-81 Effect of high p-palmitate content In infant formula on the intestinal microbiota of term Infants. *J Pediatr Gastroenterol*
39. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr*. 2013;98(2):561S-71S.
40. Vandenplas Y et al. A partly fermented infant formula with postbiotics including 3'-GL, specific oligosaccharides, 2'-FL, and milk fat supports adequate growth, is safe and well-tolerated in healthy term infants: a double-blind, randomised, controlled, multi-country trial. *Nutrients*. 2020; 12(11):3960
41. Overbeek S et al. Human milk oligosaccharide 2'-fucosyllactose more efficiently modulates immunogenicity during maturation of human dendritic cells in the presence of scGOS/lcFOS prebiotics. *JPGN*. 2019;58:995
42. Dinleyici M, Barbieur J, Dinleyici EC, Vandenplas Y. Functional effects of human milk oligosaccharides (HMOs). *Gut Microbes*. 2023 Jan-Dec;15(1):2186115.
43. Scholtens PA et al. Stool characteristics of infants receiving short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides: a review. *World J Gastroenterol*. 2014;20(37):13446-52.

Referências Fortini Complete, Fortini Plus, Fortini Plus Mf, Infatrini

1. Sociedade Brasileira de Pediatria – Departamento de Nutrologia. Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar - 4^a. ed. - São Paulo: SBP, 2018
2. Samuel T. A Narrative Review of Childhood Picky Eating and Its Relationship to Food Intakes, Nutritional Status, and Growth. *Nutrients* 2018, 10(12), 1992
3. Pesquisa realizada pela Editora Abril em parceria com a Danone Nutricia. 2020.
4. Taylor et al. Picky/fussy eating in children: Review of definitions, assessment, prevalence and dietary intakes. *Appetite* 95 (2015) 349-359.
5. Kerzner B, Milano K, MacLean WC, et al. A Practical approach to classifying and managing feeding difficulties. 2015; 135(2):344-35
6. Nogueira CA, Mello E, Filho DR, Maximino P, Fisberg M. Consenso da Associação Brasileira de Nutrologia sobre o uso de suplementos alimentares para crianças com dificuldades alimentares. *Int J Nutrol* 2018;11(supl S1):S4–S15
7. Sociedade Brasileira de Pediatria. Guia de orientações - Dificuldades alimentares. Departamento Científico de Nutrologia São Paulo: SBP, 2022.
8. Braegger C, Decsi T, Dias JA, Hartman C, Kolaček S, Koletzko B et al. Practical Approach to Paediatric Enteral Nutrition: A Comment by the ESPGHAN Committee on Nutrition. *JPGN*; 2010;51(1):110-122
9. Manual de Suporte Nutricional da Sociedade Brasileira de Pediatria - 2ed. - Rio de Janeiro: Departamento Científico de Suporte Nutricional da Sociedade Brasileira de Pediatria. – 2020.
10. NICE Guideline. Faltering growth: recognition and management of faltering growth in children. 2017.
11. Mehta NM et al. Defining pediatric malnutrition: A paradigm shift toward etiology-related definitions. *J Parenter Enter Nutr.* 2013;37(4):460–81.
- 12.. Maximino P, Ricci R, Machado RHV et al. Children with feeding difficulties have insufficient intake of vitamins, minerals, and dietary fiber. *Nutrire* volume 45, Article number: 22 (2020)
13. Xue Y, Lee E, King K et al., Prevalence of picky eating behaviour in Chinese school-age children and associations with anthropometric parameters and intelligence quotient. A cross-sectional study. *Appetite* 91 (2015) 248–255.
14. Antoniou E, Roefs A, Kremers S et al. Picky eating and child weight status development: a longitudinal study. *Journal of Human Nutrition and Dietetics.* 2015; 29(3):298-307
15. Taylor, C., & Emmett, P. Picky eating in children: Causes and consequences. *Proceedings of the Nutrition Society.* 2019; 78(2): 161-169.

16. Kreipe, R.E., Palomaki, A. Beyond Picky Eating: Avoidant/Restrictive Food Intake Disorder. *Curr Psychiatry Rep* 14, 421–431 (2012). 19.
17. Fernandez C, McCaffery H, Miller A et al. Trajectories of Picky Eating in Low-Income US Children. *Pediatrics*. 2020; 145 (6): e20192018.
18. Nogueira-de-Almeida, C.A.; Del Ciampo, L.A.; Martinez, E.Z.; Contini, A.A.; Nogueira-de-Almeida, M.E.; Ferraz, I.S.; Epifanio, M.; da Veiga Ued, F. Clinical Evolution of Preschool Picky Eater Children Receiving Oral Nutritional Supplementation during Six Months: A Prospective Controlled Clinical Trial. *Children* 2023, 10, 495.
19. Devaera, Y., Syaharutsa, D. M., Jatmiko, H. K., & Sjarif, D. R. Comparing Compliance and Efficacy of Isocaloric Oral Nutritional Supplementation Using 1.5 kcal/mL or 1 kcal/mL Sip Feeds in Mildly to Moderately Malnourished Indonesian Children: A Randomized Controlled Trial. *Pediatric gastroenterology, hepatology & nutrition*, 21(4), 315–320.
20. Food and Agriculture Organization of the United Nations. (2010). Fats and fatty acids in human nutrition. Report of an expert consultation. Food and Nutrition Paper 91.
21. Scholtens PAMJ, Goosens DAM, Staiano A. (2014). Stool characteristics of infants receiving shortchain galacto-oligosaccharides and long-chain fructooligosaccharides: A review. *World J Gastroenterol*, 20(37), 13446-13452.
22. Knol et al. (2005). Colon microflora in infants fed formula with galacto- and fructooligosaccharides: more like breast-fed infants. *J Pediatr Gastroenterol Nutr*, 40(1), 36-42
23. ILSI Brasil. Serie de publicações funções plenamente reconhecidas de nutrientes. 2019

Referências Diagnóstico E Tratamento De Aplv

1. Ed. Abril. Alergia à proteína do leite de Vaca. Revista Veja Saúde, 2020.
2. Atualização em Alergia Alimentar 2025: posicionamento conjunto da ASBAI e SBP – de Oliveira LCL, et al. *Arq Asma Alerg Imunol* – Vol. 9, Nº 1, 2025
3. Koletzko S et al. Diagnostic approach and management of cow's-milk protein allergy in infants and children: ESPGHAN GI Committee practical guidelines. *J Pediatr Gastroenterol Nutr*. 2012;55 (2):221-9.
4. Muraro A et al. EAACI food allergy and anaphylaxis guidelines: diagnosis and management of food allergy. *Allergy*. 2014;69(8):1008-25.
5. Fiocchi A et al. World Allergy Organization (WAO) Diagnosis and Rationale for Action against Cow's Milk Allergy (DRACMA) Guidelines. *Pediatr Allergy Immunol*. 2010;21 Suppl 21:1-125.
6. Venter C et al. Diagnosis and management of non-IgE-mediated cow's milk allergy in infancy: a UK primary care practical guide. *Clin Transl Allergy*. 2013;3(1):23.

7. Venter C et al. Better recognition, diagnosis and management of non-IgE-mediated cow's milk allergy in infancy: iMAP-an international interpretation of the MAP (Milk Allergy in Primary Care) guideline. *Clin Transl Allergy*. 2017;7:26.
8. Orsi M et al. Alergia a la proteína de la leche de vaca: Propuesta de Guía para el manejo de los niños con alergia a la proteína de la leche de vaca. *Arch. argent. pediatr*. 2009; 107(5): 459-467.
9. Ferreira CT et al. Alergia alimentar não-IgE mediada: formas leves e moderadas (guia prático de atualização da Sociedade Brasileira de Pediatria). São Paulo: SBP, 2022.
10. Toca MC et al. Consenso sobre el diagnóstico y el tratamiento de la alergia a las proteínas de la leche de vaca de la Sociedad Latinoamericana de Gastroenterología, Hepatología y Nutrición. *Rev Gastroenterol Méx*. 2022;87:235--250.
11. Fox A, Brown T, Walsh J, et al. An update to the Milk Allergy in Primary Care guideline. *Clin Transl Allergy*. 2019;9:40. Published 2019 Aug 12.
12. Vandenplas Y, et al. An ESPGHAN Position Paper on the Diagnosis, Management, and Prevention of Cow's Milk Allergy. *J Pediatr Gastroenterol Nutr*. 2024 Feb;78(2):386-413.
13. Heine RG. Gastroesophageal reflux disease, colic and constipation in infants with food allergy. *Curr Opin Allergy Clin Immunol*. 2006 Jun;6(3):220-5.
14. Benninga MA, Faure C, Hyman PE, et al. Childhood functional gastrointestinal disorders: neonate/toddler. *Gastroenterology* 2016;150:1443-55.
15. SBP. Distúrbios Gastrointestinais Funcionais No Lactente E Na Criança Abaixo De 4 Anos: Um Guia Para A Prática Diária. Departamento Científico De Gastroenterologia. 2022.
16. Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation and cow's milk allergy in formula-fed infants. *Acta Pædiatrica* 2015, ISSN 0803-5253
17. Morais MB, Spolidoro JV, Vieira MC, Cardoso AL, Clark O, Nishikawa A, Castro AP. Amino acid formula as a new strategy for diagnosing cow's milk allergy in infants: is it cost-effective? *J Med Econ*. 2016;19:1207-14.
18. Birch E. et al. The DIAMOND (DHA intake and measurement of neural development) study: a double-masked, randomized controlled clinical trial of the maturation of infant visual acuity as a function of the dietary level of docosahexaenoic acid' *Am J Clin Nutr.*, vol. 91, no.4, pg. 848–859, 2010.
19. Qawasmi A et al. Meta-analysis of LCPUFA supplementation of infant formula and visual acuity. *Pediatrics*. 2013;131(1):262-72.
20. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
21. Miles EA, Childs CE, Calder PC. Long-Chain Polyunsaturated Fatty Acids (LCPUFAs) and the Developing Immune System: A Narrative Review. *Nutrients*. 2021; 13: 247
22. Vandenplas Y. Lactose Intolerance. *Asia Pac J Clin Nutr*. 2015;24(Suppl 1):S9–13
23. Marten B et al. Medium-chain triglycerides. *International Dairy Journal*. 2006;16: 1374-1382.

24. Nutrologia Pediátrica: Temas da Atualidade em Nutrologia Pediátrica - 2021. / Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia. São Paulo: SBP, 2021.
25. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am.* 2013;60(1):49-74.
26. FERREIRA, CHT; BARRETO, BAP; TOFOLI, MHC; et al. Microbioma e pediatria na saúde e na doença. Barueri [SP]: Manole, 2024. 1.ed.
27. Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar/ Sociedade Brasileira de Pediatria. – 5.ed. rev. ampl.– São Paulo: SBP, 2024. 208 p.
28. Arslanoglu S, Moro GE, Boehm G, Wienz F, Stahl B, Bertino E. Early neutral prebiotic oligosaccharide supplementation reduces the incidence of some allergic manifestations in the first 5 years of life. *J Biol Regul Homeost Agents.* 2012;26(3 Suppl):49-59.
29. Arslanoglu S, Moro GE, Schmitt J, Tandoi L, Rizzardi S, Boehm G. Early Dietary intervention with a mixture of prebiotic oligosaccharides reduces the incidence of allergic manifestations and infections during the first two years of life. *J Nutr.* 2008;138(6):1091-5.
30. Boehm G. A mixture of prebiotics oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Arch Dis Child.* 2006;91(10):814
31. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5
32. Koletzko B, Baker S, Cleghorn G, et al. Global standard for the composition of infant formula: recommendations of an ESPGHAN coordinated international expert group. *J Pediatr Gastroenterol Nutr.* 2005;41(5):584-99.
33. SALMINEN, Seppo; SZAJEWSKA, Hania; KNOL, Jan. The Biotics Family in Early Life. 2. ed. Wiley, 2023.
34. Goh CY, et al. Combination of short-chain GOS and long-chain FOS 9:1 with 2FL positively impact the infant gut microbiota composition and metabolic activity in a stimulator of the human intestinal microbial ecosystem (SHIME). *J Pediatr Gastroenterol Nutr* 2019; 68 (S1): NP 114; 1159.
35. Estudo observacional, prospectivo, aberto e multicêntrico de vigilância pós-comercialização sobre aceitabilidade e tolerabilidade de uma fórmula infantil extensamente hidrolisada. Brasil. 2016: dados preliminares.
36. Epifanio, M. et al. Palatability and Sensory Perception of Infant Formulas for the Treatment of Cow's Milk Allergy According to Brazilian Mothers. *Biomed J Sci & Tech Res* 25(1)-2020. BJSTR. MS.ID.00416.
37. Eussen SRBM, et al. Theoretical Impact of Replacing Whole Cow's Milk by Young-Child Formula on Nutrient Intakes of UK Young Children: Results of a Simulation Study. *Annals of Nutrition and Metabolism*, 2015; 67(4):247-56.
38. Sociedade Brasileira de Pediatria (SBP). Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar / Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia. – 4ª. ed. - São Paulo: SBP, 2018. 172 p.

39. Institute of Medicine (US) Committee to Review Dietary Reference Intakes for Vitamin D and Calcium; Ross AC, Taylor CL, Yaktine AL, et al., editors. Dietary Reference Intakes for Calcium and Vitamin D. Washington (DC): National Academies Press (US); 2011.
40. Savilahti EM, Savilahti E. Development of natural tolerance and induced desensitization in cow's milk allergy. *Pediatr Allergy Immunol*. 2013 Mar;24(2):114-21
41. Medeiros LCS et al. Ingestão de nutrientes e estado nutricional de crianças em dieta isenta de leite de vaca e derivados. *J. Pediatr. (Rio J.)* vol.80 no.5 Porto Alegre, 2004.
42. Meyer R, De Koker C, Dziubak R, Venter C, Dominguez-Ortega G, Cutts R et al. Malnutrition in children with food allergies in the UK. *J Hum Nutr Diet*. 2014;27(3):227-35
43. Solé D et al. Consenso Brasileiro sobre Alergia Alimentar: 2018 – Parte 1 e 2. *Arq Asma Alerg Imunol*. 2018;2(1):7-82.
44. Sociedade Brasileira de Pediatria. Guia de orientações - Dificuldades alimentares. Departamento Científico de Nutrologia São Paulo: SBP, 2022.
45. Meyer R, Venter C, Fox AT, Shah N. Practical dietary management of protein energy malnutrition in young children with cow's milk protein allergy. *Pediatr Allergy Immunol*. 2012;23(4):307-14.

Referências Linha Gastro

1. Borrelli O et al. Use of a new thickened formula for treatment of symptomatic gastroesophageal reflux in infants. *Ital J Gastroenterol Hepatol*, 1997; 29:237-42.
2. Wenzl TG et al. Effects of thickened feeding on gastroesophageal reflux in infants: a placebo-controlled crossover study using intraluminal impedance. *Journal of Pediatrics*, 2003;111(4):355-359.
3. Koletzko B et al. The roles of long-chain polyunsaturated fatty acids in pregnancy, lactation and infancy: review of current knowledge and consensus recommendations. *J Perinat Med*. 2008;36(1):5-14.
4. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am*.2013;60(1):49-74. Infante
5. Salvatore, S. et al. Thickened infant formula: what to know. *Nutrition*, 2018.
6. Rosen R, Vandenplas Y, Singendonk M, et al. Pediatric Gastroesophageal Reflux Clinical Practice Guidelines: Joint Recommendations of the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition and the European Society for Pediatric Gastroenterology, Hepatology, and Nutrition. *J Pediatr Gastroenterol Nutr*. 2018;66(3):516-554.
7. Infante Pina D et al. Thickened infant formula, rheological study of the “in vitro” properties. *An Pediatr (Barc)*. 2010; 72(5):302-8.

8. Horvath A, Dziechciarz P, Szajewska H. The Effect of Thickened-Feed Interventions on Gastroesophageal Reflux in Infants: Systematic Review and Meta-analysis of Randomized, Controlled Trials. *Pediatrics* 2008;122:e1268–e1277.
9. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5
10. Braegger C et al. Supplementation of infant formula with probiotics and/or prebiotics: a systematic review and comment by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr.* 2011;52:238–225
11. Boehm G et al. Prebiotics in infant formulas. *J Clin Gastroenterol.* 2004; 38: S76-9.
12. Vandenplas Y et al. Oligosaccharides in infant formula: more evidence to validate the role of prebiotics. *Br J Nutr.* 2015;113(9):1339-44.
13. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr.* 2013;98(2):561S-71S.
14. Comparativo de tabela nutricional de produtos da mesma categoria realizado em Maio 2025.
15. Vandenplas Y et al. Algorithms for managing infant constipation, colic, regurgitation and cow's milk allergy in formula-fed infants. *Acta Pædiatrica* 2015, ISSN 0803-5253
16. Savino F et al. Reduction of crying episodes owing to infantile colic: A randomized controlled study on the efficacy of a new infant formula. *Eur J Clin Nutr* 2006, 60:1304-10.
17. Savino F et al. “Minor” feeding problems during the first months of life: effect of a partially hydrolyzed milk formula containing fructo- and galacto-oligosaccharides. *Acta Paediatr Suppl* 2003, 91:86-90.
18. Schmelzle H et al. Randomized double-blind study of the nutritional efficacy and bifidogenicity of a new infant formula containing partially hydrolyzed protein, a high beta-palmitic acid level, and nondigestible oligosaccharides. *J Pediatr Gastroenterol Nutr* 2003, 36:343-51.
19. Bongers ME et al. The clinical effect of a new infant formula in term infants with constipation: a double-blind, randomized cross-over trial.
20. Havlicekova Z et al. Beta-palmitate - a natural component of human milk in supplemental milk formulas. *Nutr J.* 2016;15(1):28.
21. Litmanovitz I et al. Reduced crying in term infants fed high beta-palmitate formula: a double-blind randomized clinical trial. *BMC Pediatr.* 2014;14:152.
22. Yao M et al. Effects of term infant formulas containing high sn-2 palmitate with and without oligofructose on stool composition, stool characteristics, and bifidogenicity. *J Pediatr Gastroenterol Nutr.* 2014 ;59(4):440-8.

23. Nowacki J et al. Stool fatty acid soaps, stool consistency and gastrointestinal tolerance in term infants fed infant formulas containing high sn-2 palmitate with or without oligofructose: a double-blind, randomized clinical trial. *Nutr J*. 2014;13:105.

24. Simakachorn N et al. Randomized, double-blind clinical trial of a lactose-free and a lactose containing formula in dietary management of acute childhood diarrhea. *J Med Assoc Thai*. 2004 Jun;87(6):641 - 9.