

REFERÊNCIAS #1 FOR GOOD REASON:

1. NielsenIQ Retail Index Evolution, CP 188 (Leites Especiais em Pó), Segmento de Alergia, MOV'25 (Out'24-Set'25), T. BRASIL INA+INFC
2. HCP TRACKER, 2024
3. Atualização em Alergia Alimentar 2025: posicionamento conjunto da ASBAI e SBP – de Oliveira LCL, et al. Arq Asma Alerg Imunol – Vol. 9, Nº 1, 2025
4. Vandenplas Y, et al. An ESPGHAN position paper on the diagnosis, management and prevention of cow's milk allergy. J Pediatr Gastroenterol Nutr. 2023 Jul 26.
5. Verwimp JJ, et al. Symptomatology and growth in infants with cow's milk protein intolerance using two different whey protein hydrolysate based formulas in a Primary Health Care setting. Eur J Clin Nutr. 1995;49 Suppl 1:S39 S48.2.
6. Giampietro PG, et al. Hypoallergenicity of an extensively hydrolyzed whey formula. Pediatr Allergy Immunol. 2001;12(2):83 86. 3.
7. SBP. Guia de orientações - Dificuldades alimentares/ Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia . São Paulo: SBP, 2022. 66
8. Meyer R, Venter C, Fox AT, Shah N. Practical dietary management of protein energy malnutrition in young children with cow's milk protein allergy. Pediatr Allergy Immunol. 2012;23(4):307-14.
9. Sampson et al (1992). Safety of an amino acid derived infant formula in children allergic to cow milk. Pediatrics, 90(3), 463 465.4.
10. Isolauri, E., et al (1995). Efficacy and safety of hydrolyzed cow milk and amino acid derived formulas in infants with cow milk allergy. The Journal of pediatrics, 127(4), 550 557.
11. Niggemann B, Binder C, Dupont C, Hadji S, Arvola T, Isolauri E. Prospective, controlled, multi-center study on the effect of an amino-acid-based formula in infants with cow's milk allergy/intolerance and atopic dermatitis. Pediatr Allergy Immunol. 2001 Apr;12(2):78-82.
12. de Boissieu D, Matarazzo P, Dupont C. Allergy to extensively hydrolyzed cow milk proteins in infants: identification and treatment with an amino acid-based formula. J Pediatr. 1997 Nov;131(5):744-7.
13. Vanderhoof JA, Murray ND, Kaufman SS, Mack DR, Antonson DL, Corkins MR, Perry D, Kruger R. Intolerance to protein hydrolysate infant formulas: an underrecognized cause of gastrointestinal symptoms in infants. J Pediatr. 1997 Nov;131(5):741-4.
14. Kelly KJ, Lazenby AJ, Rowe PC, Yardley JH, Perman JA, Sampson HA. Eosinophilic esophagitis attributed to gastroesophageal reflux: improvement with an amino acid-based formula. Gastroenterology. 1995 Nov;109(5):1503-12. doi: 10.1016/0016-5085(95)90637-1. PMID: 7557132.
15. Markowitz JE, Spergel JM, Ruchelli E, Liacouras CA. Elemental diet is an effective treatment for eosinophilic esophagitis in children and adolescents. Am J

Gastroenterol. 2003 Apr;98(4):777-82. doi: 10.1111/j.1572-0241.2003.07390.x. PMID: 12738455.

16. Moraes MB, Spolidoro JV, Vieira MC, Cardoso AL, Clark O, Nishikawa A, Castro AP. Amino acid formula as a new strategy for diagnosing cow's milk allergy in infants: is it cost-effective? J Med Econ. 2016 Dec;19(12):1207-1214.

17. Hill DJ, Heine RG, Cameron DJ, Francis DE, Bines JE. The natural history of intolerance to soy and extensively hydrolyzed formula in infants with multiple food protein intolerance. J Pediatr. 1999 Jul;135(1):118-21.

18. Informações baseadas em dados internos da Danone.

19. Danone Nutricia Research: Neocate Junior HCP and Parent experience surveys. November, 2024.

20. Estudo observacional, prospectivo, aberto e multicêntrico de vigilância pós-comercialização sobre aceitabilidade e tolerabilidade de uma fórmula infantil extensamente hidrolisada. Brasil. 2016: dados preliminares.

21. Epifanio, M. et al. Palatability and Sensory Perception of Infant Formulas for the Treatment of Cow's Milk Allergy According to Brazilian Mothers. Biomed J Sci & Tech Res 25(1)-2020. BJSTR. MS.ID.00416.

22. Estudo independente com consumidores comparando Aptamil Pepti HMO e Pregomin Pepti com outra grande marca de fórmula extensamente hidrolisada (EHF) adequada desde o nascimento para APLV leve a moderada, com n=124 pais no Brasil (2025). Dados em arquivo.

23. Nutrologia Pediátrica: Temas da Atualidade em Nutrologia Pediátrica - 2021. / Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia. São Paulo: SBP, 2021.

24. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. Pediatr Clin North Am.2013;60(1):49-74.

25. FERREIRA, CHT; BARRETO, BAP; TOFOLI, MHC; et al. Microbioma e pediatria na saúde e na doença. Barueri [SP]: Manole, 2024. 1.ed.

26. Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar/ Sociedade Brasileira de Pediatria. – 5.ed. rev. ampl.– São Paulo: SBP, 2024. 208 p.

27. Arslanoglu S, Moro GE, Boehm G, Wienz F, Stahl B, Bertino E. Early neutral prebiotic oligosaccharide supplementation reduces the incidence of some allergic manifestations in the first 5 years of life. J Biol Regul Homeost Agents. 2012;26(3 Suppl):49-59.

28. Arslanoglu S, Moro GE, Schmitt J, Tandoi L, Rizzardi S, Boehm G. Early Dietary intervention with a mixture of prebiotic oligosaccharides reduces the incidence of allergic manifestations and infections during the first two years of life. J Nutr. 2008;138(6):1091-5.

29. Boehm G. A mixture of prebiotics oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Arch Dis Child*. 2006;91(10):814-
30. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr*. 2002;34(3):291-5
31. Koletzko B, Baker S, Cleghorn G, et al. Global standard for the composition of infant formula: recommendations of an ESPGHAN coordinated international expert group. *J Pediatr Gastroenterol Nutr*. 2005;41(5):584-99.
32. SALMINEN, Seppo; SZAJEWSKA, Hania; KNOL, Jan. *The Biotics Family in Early Life*. 2. ed. Wiley, 2023.
33. Goh CY, et al. Combination of short-chain GOS and long-chain FOS 9:1 with 2FL positively impact the infant gut microbiota composition and metabolic activity in a stimulator of the human intestinal microbial ecosystem (SHIME). *J Pediatr Gastroenterol Nutr* 2019; 68 (S1): NP 114; 1159.
34. Braegger C, Decsi T, Dias JA, Hartman C, Kolaček S, Koletzko B, et al. Practical approach to paediatric enteral nutrition: A comment by the ESPGHAN committee on nutrition. *J Pediatr Gastroenterol Nutr*. 2010;51(1):110-22
35. Rodrigues VCC, Cezar TM, Abreu CLM, Sanudo A, Morais MB. A real-world cross-sectional study evaluating the role of an oral amino acid-based supplement in nutrient intake by preschoolers on a cow's milk elimination diet. *Allergol Immunopathol (Madr)*. 2024 May 1;52(3):42-52.
36. Gibson GR. Dietary modulation of the human gut microflora using the prebiotics oligofructose and inulin. *J Nutr*. 1999;129:1438S-41S.
37. Cummings JH, Macfarlane GT. Gastrointestinal effects of prebiotics. *Br J Nutr*. 2002; 87 (Suppl 2):S145-S151.
38. Veereman G. Pediatric applications of inulin and oligofructose. *J Nutr*. 2007; 137:2585S-9S.
39. Institute of Medicine (US) Committee to Review Dietary Reference Intakes for Vitamin D and Calcium. **Dietary reference intakes for calcium and vitamin D**. Ross AC, Taylor CL, Yaktine AL, Del Valle HB, editors. Washington (DC): National Academies Press (US); 2011.