

Referências – mapas mentais

Proteína Intacta

- 1.Sociedade Brasileira de Pediatria. Manual de orientação para a alimentação do lactente, do pré-escolar, do escolar, do adolescente e na escola. Departamento de Nutrologia, 4ª ed. Rio de Janeiro, 2018.
- 2.Sociedade Brasileira de Pediatria. Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar– 5.ed. rev. ampl. – São Paulo: SBP, 2024.208 p.
- 3.Zubler et al. Evidence-informed milestones for developmental surveillance tools Pediatrics. 2022;149 (3)
- 4.Moreno Villares JM, Collado MC, Larqué E, Leis Trabazo MR, Sáenz de Pipaón M, Moreno Aznar LA. Los primeros 1000 días: una oportunidad para reducir la carga de las enfermedades no transmisibles. *Nutr Hosp*. 2019;36(1):218-232. doi:10.20960/nh.02453.
- 5.Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am*. 2013;60(1):49-74.
- 6.Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
- 7.Liu L et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. *J Lipid Res*. 2014;55(3):531-9
- 8.Mun JG, Legette LL, Ikente CJ, Mitmesser SH. Choline and DHA in Maternal and Infant Nutrition: Synergistic Implications in Brain and Eye Health. *Nutrients*.2019 May 21;11(5):1125.
- 9.Salminen S, Szajewska H, Knol J. The Biotics Family in Early Life. Edited Wiley: vol. 4, 2019
- 10.Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr*. 2002;34(3):291-5
- 11.Bruzzese E et al. A formula containing galacto- and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study. *Clin Nutr*. 2009;28(2):156-61

- 12.Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr*. 2013;98(2):561S-71S.
- 13.Scholtens PA et al. Stool characteristics of infants receiving short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides: a review. *World J Gastroenterol*. 2014;20(37):13446-52.
- 14.Vandenplas Y et al. A partly fermented infant formula with postbiotics including 3'-GL, specific oligosaccharides, 2'-FL, and milk fat supports adequate growth, is safe and well-tolerated in healthy term infants: a double-blind, randomised, controlled, multi-country trial. *Nutrients*. 2020; 12(11):3960
- 15.Overbeek S et al. Human milk oligosaccharide 2'-fucosyllactose more efficiently modulates immunogenicity during maturation of human dendritic cells in the presence of scGOS/lcFOS prebiotics. *JPGN*. 2019; 58:995
- 16.Comparativo realizado entre produtos da mesma categoria, em 03/2026, utilizando informações extraídas da tabela nutricional
- 17.Luque et al. Early Programming by Protein Intake: The Effect of Protein on Adiposity Development and the Growth and Functionality of Vital Organs. *Nutrition and Metabolic Insights* 2015;8(S1) 49–56
- 18.Nogueira-de-Almeida, C. A., Falcão, M. C., Ribas Filho, D., Zorzo, R. A., Konstantyner, T., Ricci, R., Gioia, N., & Fisberg, M. (2022). Consensus of the Brazilian Association of Nutrology on Milky Feeding of Children Aged 1–5 Years Old. *International Journal of Nutrology*, 13(1), 2–16.
- 19.Hojsak, I. et al. Young Child Formula: A Position Paper by the ESPGHAN Committee on Nutrition *JPGN* 2018,66: 177-185.
- 20.Nogueira-de-Almeida, C. A., Vilanova, K. C. M., Perini, T. M., Filho, D. R. (2024). Alimentação da Criança de Zero a Cinco Anos. Manual do Departamento de Nutrologia, ABRAN.
- 21.Arrieta MC, Stiemsma LT, Amenyogbe N, Brown EM, Finlay B. The intestinal microbiome in early life: health and disease. *Front Immunol*. 2014 Sep 5;5:427. doi: 10.3389/fimmu.2014.00427. PMID: 25250028; PMCID: PMC4155789.
- 22.Bar-Yoseph F et al. Review of sn-2 palmitate oil implications for infant health. *Prostaglandins Leukot Essent Fatty Acids*. 2013;89(4):139-43
- 23.Koletzko B et al. Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *Am J Clin Nutr*. 2009; 89: 1836–45.

24. Dallas DC, Murray NM, Gan J. Proteolytic Systems in Milk: Perspectives on the Evolutionary Function within the Mammary Gland and the Infant. *J Mammary Gland Biol Neoplasia*. 2015 Dec;20(3-4):133-47.
25. Moore DR, Soeters PB. The Biological Value of Protein. *Nestle Nutr Inst Workshop Ser*. 2015;82:39-51.
26. Manary M, Callaghan M, Singh L, Briend A. Protein Quality and Growth in Malnourished Children. *Food Nutr Bull*. 2016 Mar;37 Suppl 1:S29-36.
27. Moughan PJ, Lim WXJ. Digestible indispensable amino acid score (DIAAS): 10 years on. *Front Nutr*. 2024 Jul 3;11:1389719.
28. Milani GP, Edefonti V, De Cosmi V, Bettocchi S, Mazzocchi A, Silano M, Pietrobelli A, Agostoni C. Protein and growth during the first year of life: a systematic review and meta-analysis. *Pediatr Res*. 2023 Sep;94(3):878-891.
29. Moughan PJ, Fulgoni VL 3rd, Wolfe RR. The Importance of Dietary Protein Quality in Mid- to High-Income Countries. *J Nutr*. 2024 Mar;154(3):804-814.
30. Kunz C, Lönnerdal B. Re-evaluation of the whey protein/casein ratio of human milk. *Acta Paediatr*. 1992 Feb;81(2):107-12.
31. Gridneva Z, Tie WJ, Rea A, Lai CT, Ward LC, Murray K, Hartmann PE, Geddes DT. Human Milk Casein and Whey Protein and Infant Body Composition over the First 12 Months of Lactation. *Nutrients*. 2018 Sep 19;10(9):1332.
32. Vandenplas Y, Alarcon P, Fleischer D, Hernell O, Kolacek S, Laignelet H, Lönnerdal B, Raman R, Rigo J, Salvatore S, Shamir R, Staiano A, Szajewska H, Van Goudoever HJ, von Berg A, Lee WS. Should Partial Hydrolysates Be Used as Starter Infant Formula? A Working Group Consensus. *J Pediatr Gastroenterol Nutr*. 2016 Jan;62(1):22-35.
33. Vandenplas Y, Latiff AHA, Fleischer DM, Gutiérrez-Castrellón P, Miqdady MS, Smith PK, von Berg A, Greenhawt MJ. Partially hydrolyzed formula in non-exclusively breastfed infants: A systematic review and expert consensus. *Nutrition*. 2019 Jan;57:268-274.
34. Szajewska H, Horvath A. Meta-analysis of the evidence for a partially hydrolyzed 100% whey formula for the prevention of allergic diseases. *Curr Med Res Opin*. 2010 Feb;26(2):423-37.
35. Osborn DA, Sinn JK, Jones LJ. Infant formulas containing hydrolyzed protein for prevention of allergic disease. *Cochrane Database Syst Rev*. 2018 Oct 19;10(10):CD003664.

36.Vandenplas Y, Broekaert I, Domellöf M, Indrio F, Lapillonne A, Pienar C, et al. An ESPGHAN Position Paper on the Diagnosis, Management, and Prevention of Cow's Milk Allergy. *J Pediatr Gastroenterol Nutr.* 2024;78(2):386-413.

37.Food Allergy Update 2025: joint position statement of the Brazilian Association of Allergy and Immunology and the Brazilian Society of Pediatrics. *Arq Asma Alerg Imunol.* 2025;9(1):5-96.