

Referências Bibliográficas do Fast Guide Linha Aptamil

1. Sociedade Brasileira de Pediatria – Departamento de Nutrologia Manual de Alimentação: orientações para alimentação do lactente ao adolescente, na escola, na gestante, na prevenção de doenças e segurança alimentar / Sociedade Brasileira de Pediatria. Departamento Científico de Nutrologia. – 4ª. ed. - São Paulo: SBP, 2018. 172 p.
2. Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin North Am.* 2013;60(1):49-74.
3. Koletzko B et al. Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *Am J Clin Nutr.* 2009; 89: 1836–45.
4. Falcão MC. Dinâmica da composição lipídica das fórmulas infantis e suas implicações clínicas. *BRASPEN J* 2020; 35 (3): 294-306
5. Liu L et al. Higher efficacy of dietary DHA provided as a phospholipid than as a triglyceride for brain DHA accretion in neonatal piglets. *J Lipid Res.* 2014;55(3):531-9
6. Moro G et al. Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *J Pediatr Gastroenterol Nutr.* 2002;34(3):291-5
7. Bruzzese E et al. A formula containing galacto- and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study. *Clin Nutr.* 2009;28(2):156-61
8. Oozeer R et al. Intestinal microbiology in early life: specific prebiotics can have similar functionalities as human-milk oligosaccharides. *Am J Clin Nutr.* 2013;98(2):561S-71S.
9. Scholtens PA et al. Stool characteristics of infants receiving short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides: a review. *World J Gastroenterol.* 2014;20(37):13446-52.
10. Vandenplas Y et al. A partly fermented infant formula with postbiotics including 3'-GL, specific oligosaccharides, 2'-FL, and milk fat supports adequate growth, is safe and well-tolerated in healthy term infants: a double-blind, randomised, controlled, multi-country trial. *Nutrients.* 2020; 12(11):3960
11. Overbeek S et al. Human milk oligosaccharide 2'-fucosyllactose more efficiently modulates immunogenicity during maturation of human dendritic cells in the presence of scGOS/lcFOS prebiotics. 2019;58:995